

Chianeke Okwudili

Chemical  
engineering

17/Eng01/006

Maths assignment

Assignment VI

(a) Define mathematical modelling

Mathematical modelling is defined as the act of describing a system and its process using mathematical concepts and language. It is the process of setting up a model, solving it mathematically and interpreting the result in physical or other terms.

(b) Outline two methods of obtaining mathematical model for engineering system

(i) Transition from the physical situation (physical system) to its mathematical formulation (its mathematical model)

(ii) Solution by a mathematical method

(iii) Interpreting, and <sup>the result</sup> simulating the model.

(c) A thermometer that initially reads  $10^{\circ}\text{C}$  is used to measure the temperature of a system. The temperature of the thermometer is discovered to be  $20^{\circ}\text{C}$  after 5 mins of inserting it into the system. If the actual temperature of the system is  $25^{\circ}\text{C}$ .

(i) Develop a model for the system.

(ii) Simulate the developed model for time  $t=0$  to  $t=60$  mins using a step time of 1 min with the aid of Microsoft Excel.

(iii) Obtain the dynamic response of the system with the aid of MATLAB without using `sys` command, for  $t=0$  to  $t=60$  min using a step of 1 min.

(iv) Using either the dynamic response, write the steady-state temperature of the system, and

(v) Using the developed model equation, evaluate the temperature of the thermometer as  $t \rightarrow \infty$ .

i) Let  $T(t)$  be the temperature of the system and  $T_a$  the <sup>actual</sup> temperature, by Newton's law of cooling.

$$\frac{dT}{dt} = k(T - T_a)$$

The DE is variable separable

$$\frac{dT}{(T - T_a)} = k \cdot dt$$

Integrating both sides

$$\int \frac{dT}{(T - T_a)} = \int k \cdot dt$$

$$\ln(T - T_a) = kt + C$$

$$T - T_a = e^{kt+C}$$

$$T - T_a = e^{kt} e^C$$

$C = e^C$  --- w the initial condition

$$T(t) = T_a + Ce^{kt}$$

To find  $C$ , giving the initial condition

$$T(0) = T_a + Ce^{k(0)}$$

$$10 = 25 + C$$

$$C = 10 - 25$$

$$= -15$$

$$\therefore T(t) = 25 - 15e^{kt}$$

To find  $k$

$$\text{At } t = 5 \text{ mins, } T = 20^\circ\text{C}$$

$$20 = 25 - 15e^{kt}$$

$$-5 = -15e^{kt}$$

$$\frac{1}{3} = e^{kt}$$

$$\ln \frac{1}{3} = k$$


$$k = \frac{\ln \frac{1}{3}}{5}$$

$$= -0.22$$

$$\underline{\underline{T = 25 - 15e^{-0.22t}}}$$

X(TIME)	Y=25-15*EXP(-0.22*A2)
0	10
1.5	14.216144
3	17.24722998
4.5	19.42634963
6	20.99297047
7.5	22.11925137
9	22.92896144
10.5	23.51108123
12	23.92958096
13.5	24.23045035
15	24.44675249
16.5	24.60225723
18	24.71405329
19.5	24.79442612
21	24.85220806
22.5	24.89374887
24	24.92361354
25.5	24.94508396
27	24.96051956
28.5	24.97161657
30	24.97959448
31.5	24.98532999
33	24.98945338
34.5	24.99241778
36	24.99454897
37.5	24.99608112
39	24.99718263
40.5	24.99797452
42	24.99854384
43.5	24.99895313
45	24.99924738
46.5	24.99945892
48	24.99961101
49.5	24.99972034

# GRAPH OF TEMPERATURE AGAINST TIME



## Codes

Command window

clear all

clc

close all

~~t = 0:5:10~~ t = 0:0.5:50

T = 25 - 15 \* exp(-0.22 \* t)

Tn = Subs(T)

plot(t, Tn)

label('Time (s)')


ylabel('Temperature (C)')

title('Graph of Temperature (T) against Time (t)')

grid on

grid minor.

**Graph of Temperature(C) against Time(t)**



(iv) The steady state value of the temperature is  $25^{\circ}\text{C}$  at 30 mins of the exponential approach.

(v) As  $t$  tends to infinity, the temperature approaches the steady state value which is  $25^{\circ}\text{C}$ .

(vi)  $T = 24.9$

Recall

$$T = T_{\infty} - Ce^{-kt}$$

$$24.9 = 25 - 15e^{-0.22t}$$

$$24.9 - 25 = -15e^{-0.22t}$$

$$-0.1 = -15e^{-0.22t}$$

$$\frac{-0.1}{-15} = e^{-0.22t}$$

$$\frac{1}{150}$$

$$= e^{-0.22t}$$

$$\frac{1}{150}$$

$$\ln \frac{1}{150} = -0.22t$$

$$t = \frac{-5.01}{-0.22} = \underline{\underline{22.77 \text{ min}}}$$