

Ozoemena Augustine
18/10/2027
(Chemical Eng)

The dynamic model of a body in motion performing damped forced vibration is as equation (1)

$$\frac{d^2x}{dt^2} + 5\frac{dx}{dt} + 6x = \cos t$$

Given that when $t=0$, $x=0.1$ and $\frac{dx}{dt}=0$

- using the auxiliary equation method obtain the solution of the model in form of an expression having x as a function of t
- with the aid of a matlab mfile program, plot the relationship between x and t for $0 \leq t \leq 15$ time unit using a step size of 0.01 unit and
- write the steady-state solution of the model in form of $x = k \sin(t + \alpha)$

Sol

$$c.f = m^2 + 5m + 6 = 0$$

$$(m+3)(m+2) = 0$$

$$m = -3, -2$$

$$x = Ae^{-3t} + Be^{-2t}$$

$$f(t) = \cos t$$

$$x = (\cos t + D \sin t)$$

$$\frac{dx}{dt} = -\sin t + D \cos t$$

$$\frac{d^2x}{dt^2} = -\cos t - D \sin t$$

Substitute

$$= -(\cos t - D \sin t) + 5(-\sin t + D \cos t) + 6(\cos t + D \sin t)$$

$$\cos t(-c + 50 + 6c) + \sin t(-50 - 5c + 60) = \cos t$$

$$\cos t (5c + 50) + \sin t (50 - 5c) = \cos t$$

$$5c + 50 = 1$$

$$50 - 5c = 0$$

$$100 = 1$$

$$0 = 1/10$$

find c

$$5\left(\frac{1}{10}\right) + 5c = 1$$

$$\frac{1}{2} + 5c = 1$$

$$c = 1/10$$

$$C \cdot s = x = Ae^{-3t} + Be^{-2t} + \frac{1}{10} \cos t + \frac{1}{10} \sin t$$

subs (x=0.1) t=0

$$0.1 = Ae^{-0} + Be^{-0} + \frac{1}{10} \cos 0 + \frac{1}{10} \sin 0$$

$$0.1 = A + B + 0.1$$

$$A + B = 0 \quad \text{--- eqn 1}$$

$$dx/dt = -3Ae^{-3t} - 2Be^{-2t} - \frac{1}{10} \sin t + \frac{1}{10} \cos t$$

$$\textcircled{1} \quad dx/dt = 0 \quad t=0$$

$$0 = -3A - 2B + \frac{1}{10}$$

$$3A + 2B = \frac{1}{10}$$

$$A + B = 0 \quad \times 2$$

$$3A + 2B = 0.1$$

$$2A + 2B = 0$$

$$-A = -0.1$$

$$A = 0.1$$

To find B

$$0.1 + B = 0$$

$$B = -0.1$$

$$C \cdot s = \frac{1}{10} e^{-3t} - \frac{1}{10} e^{-2t} + \frac{1}{10} \cos t + \frac{1}{10} \sin t$$

b) MATLAB CODE

command window

```
clear
```

```
clc
```

```
close all
```

```
syms x
```

```
syms t
```

```
t = 0:0.1:15
```

```
x = 0.1*exp(-2*t) + 0.1*exp(-3*t) + 0.1*cos(t) + 0.1*sin(t)
```


```
plot(t,x)
```

```
xlabel('t'), ylabel('x')
```

```
grid on
```

```
axis equal
```

Graph



(c) Steady state

$$x_{ss} = x_{t \rightarrow \infty} = 0.1 \cos t + 0.1 \sin t = k \sin(t + \alpha)$$

Recall $k \sin(t+a) = k \sin t \cos a + k \cos t \sin a$
 $x_{ss} \rightarrow \infty = 0.1 \cos t + 0.1 \sin t = k \sin t \cos a + k \cos t \sin a$

Taking the coefficients of \cos and \sin

$$0.1 = k \sin a$$

$$0.1 = k \cos a$$

Square both sides

$$k^2 \sin^2 a + k^2 \cos^2 a = \frac{1}{100} + \frac{1}{100}$$

$$k^2 (\sin^2 a + \cos^2 a) = \frac{1}{50}$$

$$k^2 = 1/50 \quad k = \sqrt{2}/10$$

$$\frac{k \sin a}{k \cos a} = \frac{0.1}{0.1} = 1 \quad \therefore a = \frac{\pi}{4}$$

$$\tan a = 1$$

$$x_{ss} = \frac{\sqrt{2}}{10} \sin(t + \pi/4)$$