



09/04/20 CHM102

Iberu Chibuzor Grace
Medicine & Surgery
191MHS01186


Assignment on Ethers




Methoxymethane




Ethoxyethane



Butoxyethane



Methoxyethane



Ethoxypropane

2) a). Physical states:-

At room temperature, ethers are colourless, neutral liquids with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.

b). Solubility:-

Ethers are less soluble in water than are the corresponding alcohols. Lower molecular weight ethers such as methoxymethane and methoxyethane are fairly soluble in water since the molecules are able to form hydrogen bonds with the water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents.

c). Density:-

Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water.

d. Boiling point


Lower molecular mass esters have a lower boiling point than the corresponding alcohols but those esters containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling point of esters tend to approximate those of hydrocarbons of same relative molecular mass from which it can be concluded that no molecules are not associated in the liquid phase as there are no ^{sufficiently} available hydrogen for association through hydrogen bonds.

e. Reactivity

Esters are inert at moderate temperatures. Their inertness at moderate temperatures leads to their wide use as reaction media. Simple esters are not found commonly in nature but the ester linkage is present in such natural products as sugar, starches and cellulose.


3) a. Partial Dehydration of Alcohols

Simple esters are manufactured from alcohols by catalytic dehydration. The alcohol in excess and concentrated tetraoxosulphate (VI) acid is heated at a carefully maintained temperature of 140°C . This process is known as continuous esterification. If excess alcohol is not used, the temperature is as high as $170-180^{\circ}\text{C}$, further dehydration to yield alkene occurs. e.g.



b. from haloalkanes and dry silver (I) oxide

This includes heating haloalkanes with dry silver (I) oxide. e.g.



- 4) Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol
- 5) Ethylene oxide is used in the preparation of carbonic anhydrase agents, plasticizers and several synthetic textiles
- 6) Ethylene oxide is used as a gaseous sterilizing agent