

Name: Osebor Ehiaghe Osasogie  
 College: Medicine and Health science  
 Department: Dentistry and dental surgery  
 Matric number: 19/mhs09/020  
 Phy102

1a. Charging by Induction:

Electric charges can be obtained on an object without touching it, by a process called electrostatic induction.

Consider a positively charged rubber rod brought near a neutral (uncharged) conducting sphere that is insulated so that there is no conducting path to ground as shown below. The repulsive force between the protons in the rod and those in the sphere causes a redistribution of charges on the sphere so that some protons move to the side of the sphere farthest away from the rod. The region of the sphere nearest the positively charged rod has an excess of negative charge because of the migration of protons away from this location. If a grounded conducting wire is then connected to the sphere, some of the protons leave the sphere and travel to the earth. If the wire to ground is then removed, the conducting sphere is left with an excess of induced negative charge.

Finally, when the rubber rod is removed from the vicinity of the sphere, the induced negatively charge remains on the ungrounded sphere and becomes uniformly distributed over the surface of the sphere.

1b.

Handwritten solution for problem 1b:

$$k = 9 \times 10^9$$

$$q_1 + q_2 = 5 \times 10^{-5} \text{ C}$$

$$F = 1 \text{ N}$$

$$d = 2 \text{ m}$$

Calculate the charge on each sphere

Recall that:  $k = 9 \times 10^9$

$$F = \frac{k q_1 q_2}{r^2}$$

$$1 = \frac{9 \times 10^9 \times (q_1 q_2 \times 5 \times 10^{-5})}{2^2}$$

$$4 = 9 \times 10^9 \times 5 \times 10^{-5} q_1 + 9 \times 10^9 q_2$$

$$4 = 4.5 \times 10^5 q_1 + 9 \times 10^9 q_2$$

$$9 \times 10^9 q_2 - 4.5 \times 10^5 q_1 = 4 = 0$$

$$q_1 = 0.0000111 \text{ C}$$

$$q_2 = 0.000038 \text{ C}$$

$$\Rightarrow q_1 = 1.1 \times 10^{-5} \text{ C}$$

$$\Rightarrow q_2 = 3.8 \times 10^{-5} \text{ C}$$

1c.

1c.  $Q_1 = Q_2 = 8 \mu\text{C}$   
 $d = 0.5 \text{ m}$   
 determine  $\Phi$  if electric field at a point P is zero

$\tan \theta = \frac{\text{opp}}{\text{adj}}$   
 $\tan \theta = \frac{1}{0.5}$   
 $\theta = \tan^{-1}(2)$   
 $\theta = 63.4^\circ$

$x^2 = 1^2 + 0.5^2$   
 $\sqrt{x^2} = \sqrt{1.25}$   
 $x = \sqrt{1.25}$   
 $x = 1.12$

$Q_2$   $d = 0.5$   $\theta = 63.4^\circ$   $Q_1 = 8 \times 10^{-6} \text{ C}$

$E_1 = \frac{k q_1}{r^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{(1.12)^2} = 5739.795918$

$E_2 = \frac{k q_2}{r^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{1.12^2} = 5739.795918$

$E_{\text{net}} = \frac{k q}{r^2} = 9 \times 10^9 \times q = 9 \times 10^9 q$

Vector	angle	x-comp	y-comp
$E_1 = 5739.795918$	$63.4^\circ$	$E_1 \cos \theta$	$E_1 \sin \theta$
$E_2 = 5739.795918$	$63.4^\circ$	$2570.045775$	$5132.262939$
$E_{\text{net}} = 9 \times 10^9 q$	$90^\circ$	$E_{\text{net}} \cos \theta = 0$	$9 \times 10^9 q$
		$x = 0$	$z_y = 10264.52568$

1c.continuation

Magnitude =  $\sqrt{(x-x)^2 + (z-y)^2}$

$E_{\text{net}} = \sqrt{(0)^2 + (10264.52568)^2}$

Since  $E = 0$

$0 = 9 \times 10^9 q + 10264.52568$

Making q the subject of formula

$q = \frac{-10264.52568}{9 \times 10^9}$

3a.

- (i) Volume charge density,
- (ii) Surface charge density,
- (iii) Linear charge density,

3b. POTENTIAL DIFFERENCE

The electric potential difference between two points in an electric field can be defined as the work done per unit charge against electrical forces when a charge is transported from one point to the other. It is measured in Volt or Joules per Coulomb . Electric potential difference is a scalar quantity.

Consider the diagram above, suppose a test charge is moved from point to point along an arbitrary path inside an electric field . The electric field exerts a force on the charge as shown in fig 3.1. To move the test charge from to at constant velocity, an external force of must act on the charge. Therefore, the elemental work done is given as:

But

Substituting equation in yields

Then total work done in moving the test charge from to is:

From the definition of electric potential difference, it follows that:  
Putting equation in yields

## SECTION B.

4a . magnetic flux is defined as the strength of the magnetic field which can be represented by line of forces. It is represented by the symbol  $\Phi$ .mathematically given as  $\Phi=B \cdot d A$

4b.

$m = 9 \times 10^{-31} \text{ kg}$   
 $r = 1.4 \times 10^{-7} \text{ m}$   
 $B = 3.5 \times 10^{-1} \text{ weber/meter}^2$   
 $\omega = \frac{v}{r} = \frac{qB}{m}$   
 $\omega = \frac{qB}{m} = \frac{1.6 \times 10^{-19} \times 3.5 \times 10^{-1}}{9 \times 10^{-31}}$   
 $\omega = 6.222222222222222 \text{ T}^{-1}$

4c. In the question we were given paramiters such as

i .mass of the electron =  $9.11 \times 10^{-31} \text{ kg}$

ii .A radius of  $1.4 \times 10^{-7} \text{ m}$

iii. magnetic field of  $3.5 \times 10^{-1} \text{ webermeter square}$

and you are asked to find the cyclotron frequency which is equal or the same thing as angular speed. it is called cyclotron frequency because it is a frequency of an accelerator called cyclotron.

Recall that angular speed is given as  $\omega =$

q Substituting we have  $\omega = \frac{1.6 \times 10^{-19} \times 3.5 \times 10^{-1}}{9.11 \times 10^{-31}}$

$$= 6.222222222222222 \text{ T}^{-1}$$

SO since cyclotron frequency is equal to angular speed the cyclotron frequency is equal to  $= 6.222222222222222 \text{ T}^{-1}$  , having a unit as  $\text{T}^{-1}$  which is equal to the unit of frequency dimensionally.

5b .Biot-savart law states that the magnetic field is directly proportional to the product permeability of free space( $\mu$ ),the current(I),the change in length, the radius and inversely proportional to square of radius ( $r^2$  ). It can be represented mathematically by where is a constant called Permeability of free space.  
The unit of is webermetre square

5b. Magnetic Field of a Straight Current Carrying Conductor




Fig 1: A section of a Straight Current Carrying Conductor  
Applying the Biot-Savart law, we find the magnitude of the field

When the length of the conductor is very great in comparison to its distance from point  $P$ , we consider it infinitely long. That is, when  $a$  is much larger than  $x$ ,

In a physical situation, we have axial symmetry about the  $y$ -axis. Thus, at all points in a circle of radius  $r$ , around the conductor, the magnitude of  $B$  is

Equation defines the magnitude of the magnetic field of flux density  $B$  near a long, straight current carrying conductor.