

**NAME: Etukudo Samuel
Ofonime**

COLLEGE: MHS

DEPT: PHARMACY

MAT NO: 19/MHS11/055

CHEMISTRY 104

ASSIGNMENT ON ETHERS

**IUPAC NAMES OF THE
FOLLOWING ORGANIC
COMPOUNDS**

1, [1] CH_3OCH_3 : Dimethyl ether or
Methoxymethane

ii, $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$: Di ethyl ether
or ethoxyethane

iii. $[\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2]_2\text{O}$: Di butyl ether

iii. $[\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2]_2\text{O}$: Di butyl ether

iv, $\text{CH}_3\text{CH}_2\text{OCH}_3$: Ethyl methyl ether

v, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$: Ethyl propyl ether

2, PROPERTIES OF ETHERS

GENERAL PROPERTIES:

Physical states: At room temperature, ethers are colorless, neutral liquids with pleasant odors. The lower aliphatic Ethers are highly flammable gases or volatile liquids

Solubility; Ethers are less soluble in water than are the corresponding alcohols. Lower molecular weight ethers such as methoxymethane and methoxy ethane are fairly soluble in water since the molecule are able to form hydrogen bonds with the water molecules but as the hydrocarbon content of the molecules increases,

form hydrogen bonds with the water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents'

Density; Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water.

Boiling point; Low molecular mass ethers have a lower boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling point of ethers tend to approximate those of hydrocarbons of same relative molecular mass from which it can be concluded that the molecules are not associated in the liquid phase as there are no suitably available hydrogen for association through hydrogen bonds,

Reactivity; Ethers are inert at

through hydrogen bonds,

Reactivity; Ethers are inert at moderate temperature. Their inertness at moderate temperatures leads to their wide use as reaction media

Simple ethers are not found commonly in nature but the ether linkage is present in such natural products as sugars, starches and cellulose.

TWO METHODS OF PREPARING ETHERS

MANUFACTURE AND PREPARATION OF ETHERS;

Partial dehydration of alcohols Simple ethers are manufactured from alcohols by catalytic dehydration. The alcohol in excess and concentrated tetra oxo sulphate(vi) acid is heated at a carefully maintained temperature of 140°C . this process is known as continuous etherification. If excess alcohol is not used, the temperature is as high as $170-180^{\circ}\text{C}$, further dehydration

to yield alkene occurs $2\text{ROH} \text{ conc}$

continuous etherification. If excess alcohol is not used, the temperature is as high as 170-180°C, further dehydration

to yield alkene occurs $2ROH \xrightarrow{\text{conc. H}_2\text{SO}_4 / 140^\circ\text{C}} R-O-R + H_2O$ Examples
 $2CH_3CH_2OH \xrightarrow{\text{conc. H}_2\text{SO}_4 / 140^\circ\text{C}} CH_3CH_2-O-CH_2CH_3 + H_2O$

Controlled catalytic hydration of olefins $2CH_3CH=CH_2 + H_2O \xrightarrow{\text{Ag}_2O}$
 $(CH_3)_2CH-O-CH(CH_3)_2$ 2-isopropoxypropane
From Haloalkanes and dry silver (I) oxide $2RX + Ag_2O \xrightarrow{\text{warm}} R-O-R + 2AgX$

$2CH_3CH_2CH_2Cl + Ag_2O \xrightarrow{\text{warm}} CH_3CH_2CH_2OCH_2CH_2CH_3 + 2AgCl$
Propoxypropane.

Uses of ethylene oxide

1, Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol.

2, Ethylene oxide is used in the

+2AgClPropoxypropane.

Uses of ethylene oxide

1, Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol.

2, Ethylene oxide is used in the preparation of nonionic emulsifying agents, plastics, plasticizers and several synthetic textiles.

3, Ethylene oxide is used as a gaseous sterilizing agent for medical equipment.

OTHERS

1, It is also used to make antifreeze, adhesives detergents and polyesters.

2. it is also used in the production of fumigants and pesticides.