



2) In the Grignard's Synthesis of alcohols, react Grignard's reagent with  
 $CH_3CH_2CH_2CH_2COCH_2CH_2CH_3$  Show the reaction step  
 $CH_3CH_2CH_2CH_2MgCl + CH_3CH_2CH_2CH_2COCH_2CH_2CH_3$   
 Propyl Magnesium chloride      Octan-4-one  
 (Grignard reagent)       $CH_3CH_2CH_2CH_3$




3) Carbohydrates Such as Starch are major group of Natural Compound  
 Malt can be made to yield ethanol but the biological process of  
 Fermentation. The biological catalyst (enzymes) found in yeast break  
 down the Carbohydrate into ethanol at 15°C. The Starch  
 containing materials include Maltases, Peptases, Cellulases and  
 Waxing with malt to 60°C for a specific period of time are  
 contained in maltose but the enzyme diastase contained in malt  
 $2(C_6H_{10}O_5)_n + nH_2O \xrightarrow{60^\circ C / \text{diastase}} nC_{12}H_{22}O_{11}$   
 maltose

The Maltose is broken down into glucose on addition of  
 yeast which contains the enzyme maltase and at a temperature  
 of 15°C




Classification based on the number of functional groups they possess. It can be classified as follows

i) Monofunctional alcohols: Monofunctional alcohols have only one hydroxyl group (-OH) present in the alcohol structure e.g.




(Ethanol)


ii) Difunctional alcohols: Difunctional alcohols are also called glycols having two hydroxyl group present in the alcohol structure e.g.



iii) Polyfunctional alcohols: Polyfunctional alcohols also called triols, but have three hydroxyl groups present in the alcohol structure e.g.






Propane-1,2,3 triol (trifunctional alcohols)




D-aldose carbohydrate

iv) Polymers containing alcohols: Polyhydric alcohols or polyols are those ones having more than three hydroxyl group e.g.




The glucose at constant temperature of  $10^{\circ}\text{C}$  is more convenient than alcohol in the enzyme zymase contained also in the yeast.




4.)

Alkenes and alkanones are reduced to primary and secondary alcohol by hydrogenation of carbon-oxygen double bond in the presence of a catalyst such as platinum, nickel, palladium.

Catalyst or with sodium tetra chloride (III) ( $\text{Ni}, \text{Pd}, \text{Pt}$ )  
 Example: Reduction of an alkenol yields an alcohol are reduced to primary alkenols' e.g.



- Reduction of an alkenone yields a secondary alcohol




Alkylalcohol: I. favour

Mechanical Engineering

19/EN9061005

1) Classification of Alkanols

- Classification based on the number of alkyl groups or hydrogen atom  
Alkanol has the general formula "R-OH", here "R" is the alkyl group  
e.g Methyl - CH<sub>3</sub>



While 'OH' is the hydroxyl group which is the main functional group of alkanols, they are classified as follows

i) Primary alkanol: They have only one alkyl group or three or two hydrogen atom attached to the atom that carries hydroxyl group e.g




Ethanol (1°)

Methanol (1°)

ii) Secondary alkanol: Secondary alkanols have two alkyl groups or one

hydrogen atom attached to the carbon that carries the hydroxyl group e.g








2-Butanol (2°)

Propan-2-ol (2°)

Butan-2-ol (2°)

2-Propanol (2°)

iii) Tertiary alkanol: Tertiary alkanol have three alkyl group and no hydrogen atom attached to the carbon atom that carries the hydroxyl group



2 Methyl Propan-2-ol (3°)

2-methyl Butan-2-ol

2 Methyl -2- Propanol

2-methyl -2- Butanol