NAME: EMENIKE ESTHER IFEOMA

MATRIC NUMBER: 19/MHS02/047

DEPARTMENT: NURSING

COURSE CODE: CHM102

- 1. Give the IUPAC names of the following organic compounds.
 - CH3OCH3- Methoxymethane
 - CH3CH2OCH2CH3- Ethoxymethane
 - (CH3CH2CH2CH2)2O-Butoxymethane
 - CH3CH2OCH3-Methoxyethane
 - CH3CH2CH2OCH2CH3- Ethoxypropane.
- 2. Discuss the properties of ethers.
 - Physical states: At room temperature, ethers are colourless, neutral liquids with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.
 - Solubility: Ethers are less soluble in water than they are the
 corresponding alcohols. Lower molecular weight ethers such as
 methoxymethane and methoxyethane are fairly soluble in water since
 the molecules are able to form hydrogen bonds with the water
 molecules but as the hydrocarbon content of the molecules increases,
 there is a rapid decline in solubility. They are miscible with the most
 organic solvents.
 - Density: Most of the simple ethers are less dense with water,
 although the density increases with increasing relative molecular mass
 and some of the aromatic ethers are in fact denser than water.

- Boiling point: Low molecular mass ethers have a lower boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling point of ethes tend to approximate those of hydrocarbons of same relative molecular mass from which it can be concluded that the molecules are not associated in the liquid phase as there are no suitably availabe hydrogen for association through hydrogen bonds.
- Reactivity: Ethers are inert at moderate temperature. Their inertness
 at moderate temperatures leads to their wide use as reaction media.
 Simple ethers are not found in commonly in nature but the ether
 linkage is present in such natural products as sugars, starches and
 cellulose.
- 3. Discuss the explicitly two methods of preparing ethers and show equations of reaction
 - Partial dehydration of alcohols: Simple ethers are manufactured from alcohols by catalytic dehydration. The alcohol in excess and concentrated tetraoxosulphate(vi)acid is heated at a carefully manitained temperature of 140°C. This process is known as continous etherification. If excess alcohol is not used, the temperature is as high as 170-180°C, further dehydration to yield alkene occurs

Controlled catalytic hydartion of olefins
 2CH3CH=CH + H2O-----→ (CH3)2CH-O-CH(CH3)2
 2-isopropoxypropane

- 4. State three uses of ethylene oxide
 - Ethylene oxide is used as an intermediate in the hydroxylic manufacture of ethylene glycol

- Ethylene oxide is used in the preparation of nonionic emulsifying agents, plastics, plasticizers and several syntheic textiles
- Ethlene oxide is used as a gaseous sterilizing agent.