

Beta oxidation is a metabolic process involving multiple steps by which fatty acid molecules are broken down to produce energy. More specifically, beta oxidation consists in breaking down long [fatty acids](#) that have been converted to acyl-CoA chains into progressively smaller fatty acyl-CoA chains. This reaction releases acetyl-CoA, FADH₂ and NADH, the three of which then enter another metabolic process called *citric acid cycle* or [Krebs cycle](#), in which ATP is produced to be used as energy. Beta oxidation goes on until two acetyl-CoA molecules are produced and the acyl-CoA chain has been completely broken down. In eukaryotic cells, beta oxidation takes place in the [mitochondria](#), whereas in prokaryotic cells, it happens in the [cytosol](#). For beta oxidation


to take place, fatty acids must first enter the cell through the cell membrane, then bind to coenzyme A (CoA), forming fatty acyl CoA and, in the case of eukaryotic cells, enter the mitochondria, where beta oxidation occurs.

Beta oxidation takes place in three steps: dehydrogenation, hydration and oxidation. Each step is catalyzed by a distinct enzyme.

Dehydrogenation


In the first step, acyl-CoA is oxidized by the enzyme acyl CoA dehydrogenase. A double bond is formed between the second and third carbons (C2 and C3) of the acyl-CoA chain entering the beta oxidation cycle; the end product of this reaction is trans- Δ^2 -enoyl-CoA (trans-delta 2-enoyl CoA).

This step uses FAD and produces FADH₂, which will enter the citric acid cycle and form ATP to be used as energy. (Notice in the following figure that the carbon count starts on the right side: the rightmost carbon below the oxygen atom is C1, then C2 on the left forming a double bond with C3, and so on.)



Hydration

In the second step, the double bond between C2 and C3 of trans- Δ^2 -enoyl-CoA is hydrated, forming the end product L- β -hydroxyacyl CoA, which has a hydroxyl group (OH) in C2, in place of the double bond. This reaction is catalyzed by another enzyme: enoyl CoA hydratase. This step requires water.



Oxidation

In the third step, the hydroxyl group in C2 of L- β -hydroxyacyl CoA is oxidized by NAD^+ in a reaction that is catalyzed by 3-hydroxyacyl-CoA dehydrogenase. The end products are β -ketoacyl CoA and $\text{NADH} + \text{H}^+$. NADH will enter the citric acid cycle and produce ATP that will be used as energy.

