## NAME; Elepo khadijah opeoluwa

## MATRIC NO; 19/mhs02/046

## **DEPARTMENT;** Nursing

1. Give the IUPAC names of the following compounds;

- HCOOH- Methanoic acid
- •HOOCCH2CH2CH2COOH-Pentan-1,5-dioic acid
- CH3CH2CH2COOH-Butanoic acid
- HO2C-CO2H-Ethanedioic acid
- CH3(CH2)4COOH-Hexanoic acid
- •CH3CH=CHCH2CH2COOH-Hex-4-eneoic acid

2. Discuss briefly the physical properties of carboxylic acids under the following headings;

•Physical appearance– All simple aliphatic carboxylic acids up to C10 are liquids at room temperature. Most other carboxylic acids are solid at room temperature although anhydrous carboxylic acid (acetic acid) also known as glacial ethanoic acid freezes to an ice-like solid below the room temperature.

•Boiling point – Boiling point increases with increasing relative molecular mass. Aromatic carboxylic acids are crystalline solids and have higher melting points than their aliphatic counterparts of comparable relative molecular mass.

• Solubility– Lower molecular mass carboxylic acids with up to four carbon atoms in their molecules are soluble in water; this largely due to their ability to form hydrogen bonds with water molecules. The water solubility of the acids decreases as the relative molecular mass increases because the structure becomes relatively more hydrocarbon in nature and hence covalent. All carboxylic acids are soluble in organic solvents.

3. Write two industrial preparations of carboxylic acids

• From Carbon(II) oxide

Methanoic acid (formic acid) is manufactured by adding carbon(II)oxide under pressure to hot aqueous solution of sodium hydroxide. The free carboxylic acid is liberated by careful reaction with tetraoxosulphate (vi) acid (H2SO4)

CO—NaOH——> HCOONa——H2SO4—> HCOOH + NaHSO4 •From petroleum

Liquid phase air oxidation of C5-C7 alkanes, obtainable from petroleum at high temperature and pressure will give C5-C7 carboxylic acids with methanoic, propanoic and butanedioic acids as by-products.

C5-C7—O2/ High temperature and pressure—>> C5-C7 carboxylic acids.

4. With equations and brief explanation discuss the synthetic preparation of carboxylic acid

•Carbonation of Grignard reagent

Aliphatic carboxylic acids are obtained by bubbling carbon (IV) oxide into the Grignard reagent and then hydrolyzed with dilute acid

RMgBr + CO2 (C2H5)2O RCOOMgBr H2O/ dil. acid RCOOH + MgBrOH R may be 10, 20, 30 aliphatic alkyl or aryl radical

In the preparation of benzoic acid, the reagent is added to solid carbon (IV) oxide (dry ice) which also serves as coolant to the reaction mixture

C6H5MgBr + CO2 (C2H5)2O C6H5COOMgBr ---H20/H---> + C6H5COOH + MgBrOH

5. With chemical equation only, outline the reduction, decarboxylation and esterification of carboxylic acid

• 4RCOOH + 3LiAlH4 (C2H5)20-> (RCH2O)4AlLi + 2LiAlO2 + 4H2 | 4H2O

4RCH2OH + Al(OH)3 + LiOH

CH3CH2CH2COOH ——LiAlH4——

Butanoic acid

> CH3CH2CH2CH2OH

Butanol • CH3CH2CH2COONa + NaOH —-fuse—>CH3CH2CH3 + Na2CO3 Kolbe synthesis 2CH3CH2COONa + 2H2O—electrolysis/aq. CH3OH—>> CH3(CH2)2CH3 +CO2 (anode) + 2NaOH + H2(cathode) • CH3CH2CH2COOH + CH3CH2CH2OH <-------> +CH3CH2CH2COO CH2CH2CH3 + H2O.