

Name:
Matrik No:
Course code:

TAKON, VICTORIA EJEN-ENOBCE
19/MHSD1405
MWT 104

1) $\int \frac{2x}{\sqrt{4x^2-1}} dx$

let $u = \sqrt{4x^2-1} = (4x^2-1)^{1/2}$

$\frac{du}{dx} = \frac{1}{2} (4x^2-1)^{-1/2}$

$\frac{dx}{du} = 2$

$\frac{dx}{dx} = 4x (4x^2-1)^{-1/2}$

$\frac{dx}{du} = \frac{du}{4x(4x^2-1)^{1/2}} = \frac{(4x^2-1)^{1/2} du}{4x}$

we have

$2 \int \frac{x}{u} dx = 2 \int \frac{x}{4x^2-1} \cdot \frac{(4x^2-1)^{1/2} du}{4x}$

$= \frac{1}{2} \int du$

$= \frac{1}{2} u + C = \frac{1}{2} \sqrt{4x^2-1} + C$

2. $\int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx$

$= \int \sin^{-1} x (1-x^2)^{-1/2} dx$

let $u = \sin^{-1} x$

$du = (1-x^2)^{-1/2} dx$

let $u = \sin^{-1} x$

$du = (1-x^2)^{-1/2} dx$

$\int u du = \frac{u^2}{2} + C$

$= \frac{(\sin^{-1} x)^2}{2} + C$

3. $\int (\tan x)^6 \sec^2 x dx$; let $u = \tan x$, $du = \sec^2 x dx$

we have;

$\int u^6 du = \frac{u^7}{7} + C$

$= \frac{(\tan x)^7}{7} + C$