

Name: Umar Abdullahi Babaraku

Matric number: 19/MHSol/421

Assignment on Ether

1. Give the IUPAC names of the following organic compounds

Answers

CH_3OCH_3 - Methoxymethane

$\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ - Ethoxyethane

$(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{O}$ - Butoxymethane

$\text{CH}_3\text{CH}_2\text{OCH}_3$ - Methoxyethane

$\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$ - Ethoxypropane

2. Properties of ethers

i) Physical states: At room temperature, ethers are colourless, neutral liquids with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.

ii) Solubility: Ethers are less soluble in water than their corresponding alcohols. Ethers with lower molecular weight are fairly soluble in water since the molecules are able to form hydrogen bonds with the water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents.

iii) Density: Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water.


iv) Boiling point: Low molecular mass ethers have a lower boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling point of ethers tend to approximate those of hydrocarbons of some relative molecular mass from which it can be concluded that the molecules are not associated in the liquid phase as there are no suitably available hydrogen ~~for~~ for association through hydrogen bonds.

v) Reactivity: Ethers are inert at moderate temperature. Their inertness at moderate temperatures leads to their wide use as reaction media.


19/MAY/21

3) Manufacture and preparation of ethers

i) Partial dehydration of alcohols: Simple ethers are manufactured from alcohols by catalytic dehydration. The alcohol in excess and concentrated tetraoxosulphate (vi) acid is heated at a carefully maintained temperature of 140°C . (this process is known as continuous etherification). If excess alcohol is not used, the temperature is as high as $170-180^{\circ}\text{C}$, further dehydration to yield alkene occurs.




Examples




ii) From haloalkanes and dry silver (i) oxide

In this method, an alkyl halide is reacted with dry silver (i) oxide which leads to the formation of alkoxyalkane.



Example



4) Uses of ethylene oxide

- i) Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol.
- ii) Ethylene oxide is used in the preparation of nonionic emulsifying agents, plastics, plasticizers and several synthetic textiles.
- iii) Ethylene oxide is used as a gaseous sterilizing agent.