

Name: Opata Clinton Chukwurah


Dept: Aeronautical & Astronautical Engineering

Matric No: 19/ENG09/019


Course Code: PHY 102

Section A

1a)



Two neutral conducting spheres are touching one another, thus allowing for the free movement of electrons between them.




when a negatively charged balloon is brought near,


Electrons move from the left sphere to the right sphere.

This leaves an unbalanced charge on the two spheres.

(positive on the left and negative on the right). In the presence of the charged balloon has induced a separation of charge.



The next step of the induction process involves separating the spheres. The right sphere is grabbed by the insulating stand and pulled away.



Result:- If a negatively-charged balloon is used to charge the left sphere by induction, then the left sphere becomes positively-charged. The right sphere becomes negatively-charged.

$$1b) k = 9 \times 10^9$$

$$q_1 \neq q_2 = 5 \times 10^{-5} \text{ C}$$

$$f = 1 \text{ N}$$

$$d = 2 \text{ m}$$

calculate the charge on each sphere,

$$\text{Recall, } k = 9 \times 10^9$$

$$f = \frac{k q_1 q_2}{r^2}$$

$$1 = \frac{9 \times 10^9 \times (q_1 q_2 \cdot 5 \times 10^{-5})}{2^2}$$

$$H = 9 \times 10^9 \times 5 \times 10^{-5} q_1 + 9 \times 10^9 q_2$$

$$H = 45 \times 10^5 q_1 + 9 \times 10^9 q_2$$


$$9 \times 10^9 q_2 = 45 \times 10^5 q_1 + H = 0$$

$$q_1 = 1.11 \times 10^{-5} \text{ C}$$

$$\approx q_2 = 3.8 \times 10^{-5} \text{ C}$$

$$1c) Q_1 = Q_2 = 8 \mu\text{C}$$

$$d = 0.5 \text{ m}$$



$$\tan \theta = \frac{\text{OPP}}{\text{ADJ}}$$

$$\tan \theta = \frac{1}{0.5} = 2$$

$$\theta = \tan^{-1} 2$$

$$\theta = 63.4^\circ$$

$$x^2 = 1^2 + 0.5^2$$

$$\sqrt{x^2} = \sqrt{1.25}$$

$$x = 1.12$$

$$E_1 = \frac{kq_1}{r_1^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{(1.12)^2} = 5739.7959$$

$$E_2 = \frac{kq_2}{r_2^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{(1.12)^2} = 5739.7959$$

$$E_y = \frac{kq}{r^2} = \frac{9 \times 10^9 \times 2}{1} = 9 \times 10^9$$

Vector	Angle	x component	y component
$E_1 = 5739.7959$	$63.4^\circ$	$-2570.0158$	$5132.2628$
$E_2 = 5739.7959$	$63.4^\circ$	$2570.0158$	$5132.2628$
$E_y = 9 \times 10^9$	$90^\circ$	$0$	$9 \times 10^9$
		$\Sigma x = 0$	$\Sigma y = 10269.5256$

$$\text{magnitude} = \sqrt{(\sum x)^2 + (\sum y)^2}$$

$$\sum = 0$$

$$0 = 9 \times 10^9 q + 10264.53568$$

by making  $q$  the subject of the formula

$$q = \frac{-10269.53568}{9 \times 10^9}$$

$$q = 1.1410502856 \times 10^{-6}$$

$$\therefore q = 1.14 \mu\text{C}$$

3i) Volume charge density,  $\rho = \frac{q}{V}$ , where  $q$  is the charge and  $V$  is the volume of distribution. The S.I unit is  $\text{Cm}^{-3}$

3ii) Surface charge density,  $\sigma = \frac{q}{A}$ , where  $q$  is the charge and  $A$  is the area of the surface. The S.I unit is  $\text{Cm}^{-2}$

3iii) Linear charge density,  $\lambda = \frac{q}{L}$  where  $q$  is the charge and  $L$  is the length over which it is distributed. The S.I unit is  $\text{Cm}^{-1}$

3b) The potential difference between point A and B.

$V_B - V_A$  is defined to be the change in potential energy of a charge  $q$  moved from A to B, is equal to the change in potential energy divided by the charge, potential difference is commonly called voltage, represented with the symbol

$$\Delta V : \Delta V = \Delta PE / q \quad \Delta V = \Delta PE / q \quad \text{and} \quad \Delta PE = q \Delta V$$

H<sub>a</sub>) Magnetic flux is defined as the strength of the magnetic H-field which can be represented by line of force. It is represented by the symbol  $\Phi$  mathematically

given as  $\Phi = B \cdot IA$

H<sub>b</sub>)  $m = 9 \times 10^{-31} \text{ kg}$

$r = 1.4 \times 10^{-2} \text{ m}$

$B = 3.5 \times 10^{-1} \text{ webermeter}^{-2}$

Cyclotron frequency = angular speed

$$\omega = \frac{v}{r} = \frac{qB}{m}$$

$$\omega = \frac{qB}{m} = \frac{1.6 \times 10^{-19} \times 3.5 \times 10^{-1}}{9 \times 10^{-31}}$$

$$\omega = 6222222222.22222 \text{ s}^{-1}$$

H<sub>c</sub>) (was asked to find the cyclotron frequency ( $\omega$  = angular speed). Cyclotron frequency being equivalent to angular speed.

$$\therefore \text{angular speed} = 6222222222.22222 \text{ s}^{-1}$$

ans.

5) Biot-Savart law states that the magnetic field is directly proportional to the product permeability of free space ( $\mu_0$ ), the current ( $i$ ) the change in length, the radius and inversely proportional to the square of radius ( $r^2$ ).