

Name: Stella Chinedu Eusebius Okoro
Department: M.B.B.S

Matric Number: 19/mhs01/164

Assignment on Ethers

1) Organic Compounds

	LUPAC Names
a CH_3OCH_3 \longrightarrow	Methoxymethane
b $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ \longrightarrow	Ethoxyethane
c $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{O}$ \longrightarrow	Butoxymethane
d $\text{CH}_3\text{CH}_2\text{OCH}_3$ \longrightarrow	Methoxyethane
e $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$ \longrightarrow	Ethoxypropane

2) Properties of Ethers:

a Physical states: At room temperature, ethers are colourless, odourless, neutral liquids with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.

b Solubility: Ethers are less soluble in water than corresponding alcohols. Lower molecular weight ethers such as methoxymethane and ethoxyethane are fairly soluble in water since the molecules are able to form hydrogen bonds with water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents.

c Density: Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water.

d Boiling Point: Low molecular mass ethers have a low boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling points of ethers tend to ~~approx~~ approximate those of hydrocarbons of same relative molecular mass from which it can be concluded that the molecules are not associated in the liquid phase as there are no suitably available hydrogen for association through hydrogen bonds.


e Reactivity: Ethers are inert at moderate temperature. Their inertness at moderate temperature leads to their wide use as reaction media.

Simple ethers are not found commonly in nature but the ether linkage is present in such natural products such as sugars, starches and cellulose.

2) Methods of Preparing Ethers with Equations


a) Partial Dehydration of Alcohols:

Simple ethers are manufactured from alcohols by catalytic dehydration. The alcohols in excess and concentrated tetraoxosulphate(VI) acid is heated at a carefully maintained temperature of 140°C . This process is known as "continuous etherification". If excess alcohol is not used, the temperature is as high as $170-180^{\circ}\text{C}$, further dehydration to yield alkene occurs.



b) From Halogenides and Dry Silver(I) Oxide:

General formula for ethers $\Rightarrow \text{R-O-R}$, R-O-Ar , Ar-O-Ar , where $\text{R} \Rightarrow$ alkyl group and $\text{Ar} \Rightarrow$ Aryl group



As said earlier here, R represents an alkyl group while X represents a halogen. The 2 atoms of the alkyl group displace the silver from its oxide to form an ether as seen above. The 2 atoms of the alkyl group attach themselves to both sides of the single oxygen atom thereby forming an ether.

A ~~Q~~ Uses of ethylene oxide:

a Ethylene oxide is used in the preparation of non-ionic emulsifying agents; plastics, plasticizers and several synthetic textiles.

b Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol.

c Ethylene oxide is used as a gaseous sterilizing agent.