

NAME:EDIGBE JOY ELOHOR

DEPARTMENT:NURSING

MATRIC NUMBER: 18/MHS02/065

COURSE CODE: PHS 202

QUESTION 4

MALE ORGASM

The orgasm phase is divided into two parts. The first, known as emission, is the stage where ejaculation is inevitable. This is immediately followed by the second stage, ejaculation, in which strong contractions of the penile muscle, anus, and perineal muscles help propel the semen from the body. During orgasm, the reward center of the brain (specifically the cerebellum, amygdala, nucleus accumbens, and ventral tegmental area) is flooded with neurochemicals, inciting the intense emotional response associated with an orgasm.

The male orgasm is a complex system involving multiple hormones, organs, and nerve pathways. The hormone testosterone, produced in the testicles, plays a central role by enhancing the sexual desire (libido) that leads to arousal, erection, and ultimately orgasm. By contrast, low testosterone not only decreases a man's energy and mood, it makes him less responsive to sexual stimuli, both physical and mental.

With that being said, a man often only requires physical stimulation to achieve arousal, while women typically need physical and mental stimulation to achieve the same.

QUESTION 1

SPERMATOGENESIS

Spermatogenesis is the process by which haploid spermatozoa develop from germ cells in the seminiferous tubules of the testis. This process starts with the mitotic division of the stem cells located close to the basement membrane of the tubules. These cells are called spermatogonial stem cells. The mitotic division of these produces two types of cells. Type A cells replenish the stem cells, and type B cells differentiate into primary spermatocytes. The primary spermatocyte divides meiotically (Meiosis I) into two secondary spermatocytes; each secondary spermatocyte divides into two equal haploid spermatids by Meiosis II. The spermatids are transformed into spermatozoa (sperm) by the process of spermiogenesis. These develop into mature

spermatozoa, also known as sperm cells. Thus, the primary spermatocyte gives rise to two cells, the secondary spermatocytes, and the two secondary spermatocytes by their subdivision produce four spermatozoa and four haploid cells.