

Name: Aliya Mariam Doolayo
 Course: CHM 102
 Department: MBBS Medicine and Surgery
 Matrix number: 19/MH501/085

- | | |
|--|----------------|
| 1. CH_3OCH_3 | Methoxymethane |
| $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ | Ethoxyethane |
| $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{O}$ | Butoxymethane |
| $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ | Methoxyethane |
| $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$ | Ethoxypropane |

2: Physical states: At room temperature, ethers are colourless, neutral liquid with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.

ii Solubility: Ethers are less soluble in water than are the corresponding alcohols. Lower molecular weight ethers such as methoxymethane and methoxyethane are fairly soluble in water since the molecules are able to form hydrogen bonds with water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents.

iii Density: Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water.

iv Boiling point: Low molecular mass ethers have a lower boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true. The boiling point of ethers tend to approximate those of hydrocarbons of same relative molecular mass from which it can be concluded that the molecules are not associated in the liquid phase as there are no suitably available hydrogen for association through hydrogen bonds.

3. Reactivity: Ethers are inert at moderate temperature. Their inertness at moderate temperatures leads to their wide use as reaction media. Simple ethers are not found commonly in nature but the ether linkage is present in such natural products as sugars, starches and cellulose.



Allyl Mercaptan Onology 19/11/2018

3. Controlled catalytic hydration of olefins



2-isopropoxypropane

ii From Haloalkanes and dry silver(I) oxide



4. It is used as an intermediate in the hydrolytic manufacture of ethylene glycol

It is used in the preparation of nonionic emulsifying agents, plastics, plasticizers and several synthetic textiles

It is used as a gaseous sterilizing agent.