

NAME :IKINI FAVOUR CATHERINE ONYINYE

MATRIC NO:17/MHS01/155



ASSIGNMENT:

1. Write an essay on cavernous system
2. Discuss walls of the nose

ANSWER:

A. WRITE AN ESSAY ON CAVERNOUS SYSTEM

The cavernous sinus within the human head is one of the dural venous sinuses[1] creating a cavity called the lateral sellar compartment bordered by the temporal bone of the skull and the sphenoid bone, lateral to the sella turcica. It's Latin name is SINUS CAVERNOSUS.



SOURCES: It takes its source from:

- middle cerebral vein,
- sphenoparietal sinus
- superior and inferior ophthalmic veins.

DRAINAGE:

It drains to the inferior and superior petrosal sinus

STRUCTURE:

The cavernous sinus is one of the dural venous sinuses of the head. It is a network of veins that sit in a cavity, approximately 1 x 2 cm in size in an adult.[2] The carotid siphon of the internal carotid artery, and cranial nerves III, IV, V (branches V₁ and V₂) and VI all pass through this blood filled space.

NEARBY STRUCTURES TO THE CAVERNOUS SINUS

Superiorly: optic tract, optic chiasma, internal carotid artery.

Inferiorly: Foramen lacerum and the junction of the body and greater wing of sphenoid bone.

Medially: Hypophysis cerebri or (pituitary gland) and sphenoidal air sinus.

Laterally: temporal lobe with uncus.

Anteriorly: superior orbital fissure and the apex of the orbit.

Posteriorly: apex of petrous temporal bone

VENOUS CONNECTIONS

The cavernous sinus receives blood from:

- Superior and inferior ophthalmic veins
- Sphenoparietal sinus
- Superficial middle cerebral veins
- Inferior cerebral veins

Blood leaves the sinus via superior and inferior petrosal sinuses as well as via the emissary veins through the foramina of the skull (mostly through foramen ovale). There are also connections with the pterygoid plexus of veins via inferior ophthalmic vein, deep facial vein and emissary veins

CONTENTS:

Apart from the blood which passes through a venous sinus, several anatomical structures, including some cranial nerves and their branches, also pass through the sinus.

Structures within the outer (lateral) wall of the compartment from superior to inferior:

- Oculomotor nerve,
- Trochlear nerve,
- Ophthalmic and maxillary branches of the trigeminal nerve,
- Structures passing through the midline (medial) wall:
- Abducens nerve,
- Internal carotid artery accompanied by the Internal carotid plexus.

These nerves, with the exception of CN V2, pass through the cavernous sinus to enter the orbital apex through the superior orbital fissure. The maxillary nerve, division V2 of the trigeminal nerve travels through the lower portion of the sinus and exits via the foramen rotundum. The maxillary branch passes external to, but immediately adjacent to, the lateral wall of the sinus.

The optic nerve lies just above and outside the cavernous sinus, superior and lateral to the pituitary gland on each side, and enters the orbital apex via the optic canal.

FUNCTIONS:

Venous drainage :

As a venous sinus, the cavernous sinus receives blood from the superior and inferior ophthalmic veins and from superficial cortical veins, and is connected to the basilar plexus of veins posteriorly. The cavernous sinus drains by two larger channels, the superior and inferior petrosal sinuses, ultimately into the internal jugular vein via the sigmoid sinus, also draining with emissary vein to pterygoid plexus.

APPLIED ANATOMY:

1. The pituitary gland lies between the two paired cavernous sinuses. An abnormally growing pituitary adenoma, sitting on the bony sella turcica, will expand in the direction of least resistance and eventually compress the cavernous sinus. Cavernous sinus syndrome may result from mass effect of these tumors and cause ophthalmoplegia (from compression of the oculomotor nerve, trochlear nerve, and abducens nerve), ophthalmic sensory loss (from compression of the ophthalmic nerve), and maxillary sensory loss

(from compression of the maxillary nerve). A complete lesion of the cavernous sinus disrupts CN III, IV, and VI, causing total ophthalmoplegia, usually accompanied by a fixed, dilated pupil. Involvement of CN V (V1 and variable involvement of V2) causes sensory loss in these divisions of the trigeminal nerve. Horner's syndrome can also occur due to involvement of the carotid ocular sympathetics, but may be difficult to appreciate in the setting of a complete third nerve injury.

2. It is the only anatomic location in the body in which an artery travels completely through a venous structure. If the internal carotid artery ruptures within the cavernous sinus, an *arteriovenous fistula* is created (more specifically, a carotid-cavernous fistula). Lesions affecting the cavernous sinus may affect isolated nerves or all the nerves traversing through it.


B) DISCUSS WALLS OF THE NOSE

The human nose is the most protruding part of the face. It bears the nostrils and is the first organ of the respiratory system. It is also the principal organ in the olfactory system. The shape of the nose is determined by the nasal bones and the nasal cartilages, including the nasal septum which separates the nostrils and divides the nasal cavity into two.

The main function of the nose is breathing, and the nasal mucosa lining the nasal cavity and the paranasal sinuses carries out the necessary conditioning of inhaled air by warming and moistening it. Another major function of the nose is olfaction, the sense of smell. The area of olfactory epithelium, in the upper nasal cavity, contains specialised olfactory cells responsible for this function.

The walls of the nasal cavity are divided into two; medial and lateral walls

The medial wall : formed by the nasal septum




**MEDIAL WALL OF NASAL CAVITY
(NASAL SEPTUM)**

The lateral walls :

are irregular owing to three bony plates, the nasal conchae, which project inferiorly, somewhat like louvers

The Lateral Walls of Nasal Cavity



Features on the lateral wall of the nasal cavity

There is the presence of nasal conchae and they curve inferomedially

The nasal conchae include;

- Superior nasal concha
- middle nasal concha
- inferior nasal concha

The conchae or turbinates of many mammals (especially running mammals and those existing in extreme environments) are highly convoluted, scroll-like structures that offer a vast surface area for heat exchange. Underneath each concha in both humans with simple nasal conchae and animals with complex turbinates is a recess or meatus {passage(s) in the nasal cavity}

External nose:

The external nose presents a root (or bridge), a dorsum, and a free tip or apex. The two inferior openings are the nostrils (or nares), bounded laterally by the ala and medially by the nasal septum. The superior part of the nose is supported by the nasal, frontal, and maxillary bones; the inferior part includes several cartilages. The continuous free margin of the nasal bones and maxillae in a dried skull is termed the piriform aperture.

Nasal cavity:

The nasal cavity extends in an antero-posterior direction from the nostrils, or nares, to the choanae. The choanae are the posterior apertures of the nose. Each choana is bounded medially by the vomer, inferiorly by the horizontal plate of the palatine bone, laterally by the medial pterygoid plate, and superiorly by the body of the sphenoid bone. Posteriorly, the nasal cavity communicates with the nasopharynx, which in many respects may be regarded as the posterior portion of the cavity. The nasal cavity is related to the anterior and middle cranial fossae, orbit, and paranasal sinuses and is separated from the oral cavity by the hard palate. In addition to the nostrils and choanae, the nasal cavity presents openings for the paranasal sinuses and the nasolacrimal duct. Further openings, covered by mucosa in vivo, are found in a dried skull, e.g., the sphenopalatine foramen. The nasal cavity is divided into right and left halves (each of which may be termed a nasal cavity) by the nasal septum. Each half has a roof, floor, and medial and lateral walls.

The roof:

The roof of the nasal cavity is formed by nasal cartilages and several bones, chiefly the nasal and frontal bones, the cribriform plate of the ethmoid, and the body of the sphenoid. The floor, wider than the roof, is formed by the palatine process of the maxilla and the horizontal plate of the palatine bone, i.e., by the palate. The medial wall, or nasal septum, is formed (from anterior to posterior) by (1) the septal cartilage (destroyed in a dried skull), (2) the perpendicular plate of the ethmoid bone, and (3) the vomer. It is usually deviated to one side. The lowest part of the septum (the columella) is membranous and mobile.

Floor:

Formed by the maxilla and the palatine bones. The incisive foramen transmits branches of the sphenopalatine artery and the nasopalatine nerve for general sensation from the nasal cavity and palate.

Sphenoethmoidal recess:

The space between the superior nasal concha and the sphenoid bone, with openings from the sphenoid sinus.

Superior meatus: The space inferior to the superior nasal concha, with openings from the posterior ethmoidal air cells.

Middle meatus: The space inferior to the middle nasal concha, with openings for the frontal sinus via the nasofrontal duct, the middle ethmoidal air cells on the ethmoidal bulla, and the anterior ethmoidal air cells and maxillary sinus in the hiatus semilunaris.

Inferior meatus: The space inferior to the inferior nasal concha, with an opening for the nasolacrimal duct, which drains tears from the eye into the nasal cavity. Sphenopalatine foramen. An opening posterior to the middle nasal concha receives the nasopalatine nerve and the sphenopalatine artery from the pterygopalatine fossa into the nasal cavity.