NAME: OPAW OYE SHARON ABISOLA

MATRIC NUMBER: 19/MHS02/102

DEPARTMENT: NURSING

COURSE CODE:CHM 102

- 1 Give the IUPAC names of the following compounds
 - HCOOH- Methanoic acid
 - HOOCCH2CH2CH2COOH- Pentan-15-dioic acid
 - CH3CH2CHCOOH- Butanoic acid
 - HO2C-CO2H- Ethanedioic acid
 - CH3(CH2)4COOH-Hexanoic acid
 - CH3CH=CHCH2CH2COOH- Hex-4-eneoic acid
- 2. Discuss briefly the physical properties of carboxylic acids under the following headings
 - Physical apperance: All simples aliphatic carboxylic acids up to C10
 are liquids at room temperarure. Most other carboxylic acids are
 solid at room temperature although annhydrous carboxylic acid
 (acetic acid) also known as glacial ethanoic acid freezes to an ice
 like solid below the room temperature.
 - Boiling points: Boiling points increases with increasing relative molecular mass. Aromatic carboxylic acids are crystalline solids have higher melting points than their aliphatic counterparts of comparable relative molecular mass.
 - Solubility: Lower molecular mass carboxylic acids with up to four

carbon atoms in their molecules are soluble in water; this largely due to their ability to from hydrogen bonds with water molecules. The water solubility of the acids decreases as the relative molecular mass increases because the structure becomes relatively more hydrocarbon in nature and hence covalent. All carboxylic acids are soluble in organic solvents.

- 3. Write two industrial preparations of carboxylic acids.
 - From carbon(ii) oxide: Methanoic acid is manufactured by adding carbon(ii)oxide under pressure to hot aqueous solution of sodium hydroxide. The free carboxylic acid is liberated by careful reaction with tetraoxosulphate(Vi) acid (H2SO4)

NaOH H2SO4
CO-----→ HCOONa------→ HCOOH+NaHSO4

 From Ethanol: Ethanoic acid is obtained commercially by the liquid phase air-oxidation of 5% solution of ethanal to ethanoic acid using manganite(ii) ethanoate catalyst. Ethanal itself is obtained from ethylene.

- 4. With equations and brief explanations discuss the synthetic preparation of carboxylic acid
 - Oxidation of primary alcohols and aldehydes: oxidation of primary alohols and aldehydes can be used to prepare carboxylic acids using the usual oxidizing agents (i.e k2cr207 or Kmn04) in acidic solution

[0] excess acid/KM nO4 [0]

RCH20H-----→RCH0-----→RCOOH

• Carbonation of Gringard reagent: Aliphatic carboxylic acids are obtained by bubbling carbon(IV) oxide into the Gringnard reagent and then hydrolyzed with dilute acid.

(C2H3)2O H2O/Dil. acid RMgBr +CO2-----→RCOOMgBr------→RCOOH+MgBrOH • Hydrolysis of nitriles(cyanides) or esters **H**+ RCN+2H2O-----→RCOOH+NH4+ H2O/H+reflux H+ H2O/H+reflux CH3CH2COOCH3------→CH3CH2COOH+CH3OH 5. With chemical equation only, outline the reduction, decarboxylation and esterification of carboxylic acid. I 4H20 4RCH2OH+AI(OH) +LiOH

LiA IH4

С	H3CH2CH2COOH→CH3CH2CH2CH2OH
В	utanoic acid butanol
•	D ecarboxylation
	fuse
	CH₃CH2CH2COONa+NaOH
	Kolbe synthesis
	Electrolysis/aq. CH30H
	2CH3CH2COONa+2H2O
	→CH3(CH2) ₂ CH3+CO _{2(Anode)} +2NaOH+H _{2(cathode)}

• Estherification

 H^{+}

CH3CH2CH2COOH+CH3CH2CH2OH←→CH3CH2CH2COOCH2CH2CH3+H2O