

NAME: UBAH CHIGOZIRIM DIVINE FAVOUR

MATRIC NO: 18/MHS01/350

DEPARTMENT: MBBS

LEVEL: 200

COLLEGE: MHS

COURSE CODE: ICBS

DEVELOPMENT

• DISCUSS THE 2ND WEEK OF EMBRYONIC DEVELOPMENT

The second week comprises of days 8-14 and the events that occur include;

- Completion of implantation of blastocysts.
- Formation of bilaminar embryonic disc (epiblast and hypoblast).
- Formation of extra embryonic structures.

• Completion of implantation of blastocysts:

As the blastocyst embeds itself in the endometrium it differentiates into two layers: the **cytotrophoblast** (inner) and **syncytiotrophoblast** (outer). The syncytiotrophoblast invades into the maternal endometrium, and in this sense it is more invasive than any tumor tissue. As it comes into contact with blood vessels it creates lacunae, or spaces which fill with maternal blood. These lacunae fuse to form lacunar networks. The maternal blood that flows in and out of these networks exchanges nutrients and waste products with the fetus, forming the basis of a primitive uteroplacental circulation.

Syncytiotrophoblast

The syncytiotrophoblast is acellular and does not expand mitotically. The syncytiotrophoblast produces human chorionic gonadotrophin (hCG), a glycoprotein hormone that stimulates the production of progesterone by the corpus luteum.

Cytotrophoblast

The cytotrophoblast is cellular and expands mitotically into the syncytiotrophoblast to form primary chorionic villi. Cells from these villi can be removed for early genetic testing at some risk to the fetus (chorionic villus sampling).

• Formation of bilaminar embryonic disc (epiblast and hypoblast):

After implantation, the inner cell mass subdivides into a bilaminar disc consisting of the **hypoblast** and **epiblast**.

Hypoblast

Hypoblast cells migrate along the inner surface of the cytotrophoblast and will form the primary yolk sac. The primary yolk sac becomes reduced in size and is known as the secondary yolk sac. In humans the yolk sac contains no yolk but is important for the transfer of nutrients between the fetus and mother.

Epiblast

Epiblast cells cavitate to form the amnion, an extra-embryonic epithelial membrane covering the embryo and amniotic cavity. Cells from the epiblast will also eventually form the body of the embryo.

- Formation of extra-embryonic structures:

Extra-embryonic mesoderm cells migrate between the cytotrophoblast and yolk sac and amnion. Extraembryonic somatic mesoderm lines the cytotrophoblast and covers the amnion. Extraembryonic somatic mesoderm also forms the connecting stalk that is the primordium of the umbilical cord. Extraembryonic visceral mesoderm covers the yolk sac. At the end of the second week it is possible to distinguish the dorsal (amniotic cavity) from the ventral (yolk sac) side of the embryo.

HIGHLIGHTS of 2nd Week of Development

- **Bilaminar germ disc**
- **complete implantation on uterine wall**
- **Trophoblast** differentiated in 2 layers
- **2 cavities formed-** amniotic cavity, extraembryonic coelomic cavity (chorionic cavity)
- **2 layers of extraembryonic mesoderm-** somato & splanchnopleure
- **Beginning of uteroplacental circulation**

