

Name Olayole Boluwatife Emmanuel
Course CHM 102
Department Medicine and Surgery
Matric No 19/MH501/343

Assignment

- ① Give the IUPAC names of the following compounds
- ① HCOOH — Methanoic acid
 - ② $\text{HOOCCH}_2\text{CH}_2\text{CH}_2\text{COOH}$ — Pentan-1,5-dioic acid
 - ③ $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$ — Butanoic acid
 - ④ $\text{HO}_2\text{C}-\text{CO}_2\text{H}$ — Ethanedioic acid
 - ⑤ $\text{CH}_3(\text{CH}_2)_4\text{COOH}$ — Hexanoic acid
 - ⑥ $\text{CH}_3\text{CH}=\text{CHCH}_2\text{CH}_2\text{COOH}$ — Hex-4-enoic acid
- ② Discuss briefly the physical properties of carboxylic acids under the following headings
- ① Physical appearance
 - ② Boiling point
 - ③ Solubility

Physical appearance:

All simple aliphatic carboxylic acid up to C_{10} are liquids at room temperature. Most other carboxylic acids are solid at room temperature although anhydrous carboxylic acid (acetic acid) also known as glacial ethanoic acid freezes to an ice-like solid below the room temperature.

Boiling point:

Boiling point increases with increasing relative molecular mass. Aromatic carboxylic acids are crystalline solids and have higher melting points than their aliphatic counterparts of comparable relative molecular mass.


Solubility:

Lower molecular carboxylic acids with up to four carbon atoms in their molecules are soluble in water; this largely due to their ability to form hydrogen bonds with water molecules. The water solubility of the acids decreases as the relative molecular mass increases because the structure become relatively more hydrocarbon in nature and hence covalent. All carboxylic acids are soluble in organic solvents.

③ Write two industrial preparation of carboxylic acids


① From Carbon (II) oxide

Methanoic acid (formic acid) is manufactured by adding carbon (II) oxide under pressure to hot aqueous solution of sodium hydroxide. The free carboxylic acid is liberated by careful reaction with tetraoxosulphate (VI) acid (H_2SO_4)



② From ethanal


Ethanoic acid is obtained commercially by the liquid phase air-oxidation of 5% solution of ethanal to ethanoic acid using manganite (II) ethanoate catalyst. Ethanal itself is obtained from ethylene.



④ With equations and brief explanation discuss the synthetic preparation of Carboxylic acid.





① Oxidation of Primary alcohol and aldehydes

Oxidation of primary alcohols and aldehydes can be used to prepare carboxylic acids using the usual oxidizing agents (i.e. $K_2Cr_2O_7$ or $KMnO_4$) in acidic solution.



⑤ With chemical equations only, outline the reduction, decarboxylation and esterification of Carboxylic acid.


~~Production~~ Reduction




Butanoic acid

Butanol

- Decarboxylation



Kolbe synthesis



- Esterification

