

## Physiology assignment 2

Bara Marion .A

18/MHS02/053

### Cyclic changes in breasts

Each month, women go through changes in

the hormones that make up the normal menstrual cycle. The hormone estrogen is produced by the ovaries in the first half of the menstrual cycle. It stimulates the growth of milk ducts in the breasts. The increasing level of estrogen leads to ovulation halfway through the cycle. Next, the hormone progesterone takes over in the second half of the cycle. It stimulates the formation of the milk glands. These hormones are believed to be responsible for the cyclical changes that many women feel in their breasts just before menstruation. These include swelling, pain, and soreness.

During menstruation, many women also have changes in breast texture. Their breasts may feel very lumpy. This is because the glands in the breast are enlarging to get ready for a possible pregnancy. If pregnancy does not happen, the breasts go back to normal size. Once menstruation starts, the cycle begins again.

What happens to the breasts during pregnancy and milk production?

Many healthcare providers believe the breasts are not fully mature until a woman has given birth and made milk. Breast changes are one of the earliest signs of pregnancy. This is a result of the hormone progesterone. In addition, the dark areas of skin around the nipples (the areolas) begin to swell. This is followed by the rapid swelling of the breasts themselves. Most pregnant women feel soreness down the sides of the breasts, and nipple tingling or soreness. This is because of the growth of the milk duct system and the formation of many more lobules.

By the fifth or sixth month of pregnancy, the breasts are fully capable of producing milk. As in puberty, estrogen controls the growth of the ducts, and progesterone controls the growth of the glandular buds. Many other hormones also play vital roles in milk production. These include follicle-stimulating hormone (FSH), luteinizing hormone (LH), prolactin, oxytocin, and human placental lactogen (HPL).

Other physical changes happen as well. These include the blood vessels in the breast becoming more visible and the areola getting larger and darker. All of these changes are in preparation for breastfeeding the baby after birth.

What happens to the breasts at menopause?

By the time a woman reaches her late 40s and early 50s, perimenopause is starting or is well underway. At this time, the levels of estrogen and progesterone begin to change. Estrogen levels dramatically decrease. This leads to many of the symptoms commonly linked to menopause. Without estrogen, the breast's connective tissue becomes dehydrated and is no longer elastic. The breast tissue, which was prepared to make milk, shrinks and loses shape. This leads to the "saggy" breasts associated with women of this age.

Women who are taking hormone therapy may have some of the premenstrual breast symptoms that they had while they were still menstruating, such as soreness and swelling. But if a woman's breasts were saggy before menopause, this will not change with hormone therapy.

### Cyclic changes in cervix

- Mucosa of uterine cervix does not undergo cyclic desquamation as the body of uterus
- regular changes in cervical mucus under influence: 1. estrogens (ovulation) mucus

thinner and more alkaline - promotion of survival and transport of sperm, spinnbarkeit at ovulation

2. progesterone (after ovulation, during pregnancy) mucus thick, tenacious, cellular

2

## Hormones and the menstrual cycle

The menstrual cycle is complex and is controlled by many different glands and the hormones that these glands produce. A brain structure called the hypothalamus causes the nearby pituitary gland to produce certain chemicals, which prompt the ovaries to produce the sex hormones oestrogen and progesterone.

The menstrual cycle is a biofeedback system, which means each structure and gland is affected by the activity of the others.

Back to top

## Phases of the menstrual cycle

The four main phases of the menstrual cycle are:

- menstruation
- the follicular phase
- ovulation
- the luteal phase.

## Menstruation

Menstruation is the elimination of the thickened lining of the uterus (endometrium) from the body through the vagina. Menstrual fluid contains blood, cells from the lining of the uterus (endometrial cells) and mucus. The average length of a period is between three days and one week.

Sanitary pads or tampons are used to absorb the menstrual flow. Both pads and tampons need to be changed regularly (at least every four hours). Using tampons has been associated with an increased risk of a rare illness called toxic shock syndrome

## Follicular phase

The follicular phase starts on the first day of menstruation and ends with ovulation. Prompted by the hypothalamus, the pituitary gland releases follicle stimulating hormone (FSH). This hormone stimulates the ovary to produce around five to 20 follicles (tiny nodules or cysts), which bead on the surface.

Each follicle houses an immature egg. Usually, only one follicle will mature into an egg, while the others die. This can occur around day 10 of a 28-day cycle. The growth of the follicles stimulates the lining of the uterus to thicken in preparation for possible pregnancy.

## Ovulation

Ovulation is the release of a mature egg from the surface of the ovary. This usually occurs mid-cycle, around two weeks or so before menstruation starts.

During the follicular phase, the developing follicle causes a rise in the level of oestrogen. The hypothalamus in the brain recognises these rising levels and releases a chemical called gonadotrophin-releasing hormone (GnRH). This hormone prompts the pituitary gland to produce raised levels of luteinising hormone (LH) and FSH.

Within two days, ovulation is triggered by the high levels of LH. The egg is funnelled into the fallopian tube and toward the uterus by waves of small, hair-like projections. The life span of the typical egg is only around 24 hours. Unless it meets a sperm during this time, it will die.

When you want to have a baby you can improve your chance of getting pregnant if you know about

ovulation and the 'fertile window' in the menstrual cycle and ovulation and menstrual syndrome

Luteal phase

During ovulation, the egg bursts from its follicle, but the ruptured follicle stays on the surface of the ovary. For the next two weeks or so, the follicle transforms into a structure known as the corpus luteum. This structure starts releasing progesterone, along with small amounts of oestrogen. This combination of hormones maintains the thickened lining of the uterus, waiting for a fertilised egg to stick (implant).

If a fertilised egg implants in the lining of the uterus, it produces the hormones that are necessary to maintain the corpus luteum. This includes human chorionic gonadotrophin (HCG), the hormone that is detected in a urine test for pregnancy. The corpus luteum keeps producing the raised levels of progesterone that are needed to maintain the thickened lining of the uterus.

If pregnancy does not occur, the corpus luteum withers and dies, usually around day 22 in a 28-day cycle. The drop in progesterone levels causes the lining of the uterus to fall away. This is known as menstruation.