

NAME - MARKMANUEL AGBEZIBE MARKBERE WINNER

DEPT - MECHANICAL ENGINEERING


MATRIC NO – 19/ENG06/035

CHEM 102: GENERAL CHEMISTRY II

Question

Assignment

1. Give the IUPAC names of the following organic compounds



2. Discuss the properties of ethers

3. Discuss explicitly two methods of preparing ethers and show equations of reaction

4. State three uses of ethylene oxide

ANSWERS

- CH_3OCH_3 - Methoxymethane
 $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ - Ethoxyethane
 $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{O}$ - Butoxymethane
 $\text{CH}_3\text{CH}_2\text{OCH}_3$ - Methoxyethane
 $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$ - Ethoxypropane

2. Properties of ethers

I. Physical states

At room temperature, ethers are colourless, neutral liquids with pleasant odours. The lower aliphatic ethers are highly flammable gases or volatile liquids.

II. Solubility

Ethers are less soluble in water than are the corresponding alcohols. Lower molecular weight ethers such as methoxymethane and methoxyethane are fairly soluble in water since the molecules are able to

form hydrogen bonds with the water molecules but as the hydrocarbon content of the molecules increases, there is a rapid decline in solubility. They are miscible with most organic solvents.

III. Density

Most of the simple ethers are less dense than water, although the density increases with increasing relative molecular mass and some of the aromatic ethers are in fact denser than water

IV. Boiling point

Low molecular mass ethers have a lower boiling point than the corresponding alcohols but those ethers containing alkyl radicals larger than four carbon atoms, the reverse is true.


V. Reactivity

Ethers are inert at moderate temperature. Their inertness at moderate temperatures leads to their wide use as reaction media.

3.

I. Partial dehydration of alcohols


Simple ethers are manufactured from alcohols by catalytic dehydration. The alcohol in excess and concentrated tetraoxosulphate(vi) acid is heated at a carefully maintained temperature of 140°C. This process is known as continuous etherification. If excess alcohol is not used, the temperature is as high as 170-180°C, further dehydration to yield alkene occurs.



Examples



II. From Haloalkanes and dry silver (I) oxide



Propoxypropane

4. Uses of ethylene oxide

- I. Ethylene oxide is used as an intermediate in the hydrolytic manufacture of ethylene glycol
- II. Ethylene oxide is used in the preparation of nonionic emulsifying agents, plastics, plasticizers and several synthetic textiles
- III. Ethylene oxide is used as a gaseous sterilizing agent