

NAME : ARIKEKPAR AZIBAOLA ISRAEL

DEPT : AERONAUTICAL & ASTRONAUTICAL ENGINEERING

MATRIC NO : 19/ENG 09/003

Name: Opata Clinton Chukwurah


Dept: Aeronautical & Astronautical Engineering

Matric No: 19/ENIG09/019


Course Code: PHY 102

Section A

1a)




Two neutral conducting spheres are touching one another, thus allowing for free movement of electrons between them.




when a negatively charged balloon is brought near,

- Electrons move from the left sphere to the right sphere.
- This leaves an unbalanced charge on the two spheres (positive on the left and negative on the right) in the presence of the charged balloon has induced a separation of charge.



The next step of the induction process involves separating the spheres. The right sphere is grounded by the metal tray stand and pulled away.



Result:- If a negatively-charged balloon is used to charge the left sphere by induction, then the left sphere becomes positively-charged. The right sphere becomes negatively-charged.

$$1b) k = 9 \times 10^9$$

$$q_1 \text{ and } q_2 = 5 \times 10^{-5} \text{ C}$$

$$f = 1 \text{ N}$$

$$d = 2 \text{ m}$$

calculate the charge on each sphere

$$\text{Recall, } k = 9 \times 10^9$$

$$f = \frac{k q_1 q_2}{r^2}$$

$$1 = \frac{9 \times 10^9 \times (q_1 \cdot 5 \times 10^{-5})}{2^2}$$

$$4 = 9 \times 10^9 \times 5 \times 10^{-5} q_1 + 9 \times 10^9 q_2$$

$$4 = 45 \times 10^5 q_1 + 9 \times 10^9 q_2$$


$$9 \times 10^9 q_2 = 45 \times 10^5 q_1 + 4 = 0$$

$$q_1 = 1.11 \times 10^{-5} \text{ C}$$

$$q_2 = 3.8 \times 10^{-5} \text{ C}$$

$$1c) Q_1 = Q_2 = 8 \mu\text{C}$$

$$d = 0.5 \text{ m}$$



$$\tan \theta = \frac{\text{OPP}}{\text{ADJ}}$$

$$\tan \theta = \frac{1}{0.5} = 2$$

$$\theta = \tan^{-1} 2$$

$$\theta = 63.4^\circ$$

$$x^2 = 1^2 + 0.5^2$$

$$\sqrt{x^2} = \sqrt{1.25}$$

$$x = 1.12$$

$$E_1 = \frac{kq_1}{r^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{(1.12)^2} = 5739.7959$$

$$E_2 = \frac{kq_2}{r^2} = \frac{9 \times 10^9 \times 8 \times 10^{-6}}{(1.12)^2} = 5739.7959$$

$$E_y = \frac{kq}{r^2} = \frac{9 \times 10^9 \times 2}{1} = 9 \times 10^9$$

Vector	Angle	x component	y component
$E_1 = 5739.7959$	$63.4^\circ$	$-2570.0158$	$5132.2628$
$E_2 = 5739.7959$	$63.4^\circ$	$2570.0158$	$5132.2628$
$E_y = 9 \times 10^9$	$90^\circ$	0	$9 \times 10^9$
		$E_x = 0$	$E_y = 10264.52568$

$$\text{Magnitude} = \sqrt{(E_x)^2 + (E_y)^2}$$

$$E = 0$$

$$0 = 9 \times 10^9 + 10264.52568$$

by making q the subject of the formula

$$q = \frac{-10264.52568}{9 \times 10^9}$$

$$q = -1.140502856 \times 10^{-6}$$

$$L. q = 1.14 \mu\text{C}$$

3i) Volume charge density,  $\rho = \frac{q}{V}$ , where  $q$  is the charge and  $V$  is the volume of distribution. The S.I unit is  $\text{Cm}^{-3}$

3ii) Surface charge density,  $\sigma = \frac{q}{A}$ , where  $q$  is the charge and  $A$  is the area of the surface. The S.I unit is  $\text{Cm}^{-2}$

3iii) Linear charge density,  $\lambda = \frac{q}{l}$  where  $q$  is the charge and  $l$  is the length over which it is distributed. The S.I unit is  $\text{Cm}^{-1}$

3b) The potential difference between point A and B,  $V_B - V_A$  is defined to be the change in potential energy of a charge  $q$  moved from A to B, is equal to the change in potential energy divided by the charge, potential difference is commonly called voltage, represented with the symbol

$$\Delta V: \Delta V = \frac{\Delta PE}{q} \quad \Delta V = \frac{\Delta PE}{q} \quad \text{and } \Delta PE = q \Delta V$$

H<sub>a</sub>) Magnetic flux is defined as the strength of the magnetic field which can be represented by the lines of force. It is represented by the symbol  $\Phi$  mathematically,

$$\text{given as } \Phi = B \cdot JA$$

$$\text{H<sub>b</sub>) } m = 9 \times 10^{-31} \text{ kg}$$

$$r = 1.4 \times 10^{-7} \text{ m}$$

$$B = 3.5 \times 10^{-1} \text{ webermeter}^{-2}$$

Cyclotron frequency = angular speed

$$\omega = \frac{v}{r} = \frac{qB}{m}$$

$$\omega = \frac{qB}{m} = \frac{1.6 \times 10^{-19} \times 3.5 \times 10^{-1}}{9 \times 10^{-31}}$$

$$\omega = 6.222222222 \cdot 222222 \cdot 10^{-1}$$

H<sub>c</sub>) (was asked to find the cyclotron frequency (i.e. angular speed). Cyclotron frequency being equivalent to angular speed.

$$\therefore \text{angular speed} = 6.222222222 \cdot 222222 \cdot 10^{-1}$$

ans.

5) Biot-Savart law states that the magnetic field is directly proportional to the product permeability of free space ( $\mu_0$ ), the current ( $i$ ), the change in length, the radius and inversely proportional to the square of radius ( $r^2$ ).