

**ORAKPO MIRABEL**  
**18/SCIO1/074**

$$\text{Let } A = \begin{Bmatrix} 3 & 1 & 2 \\ 1 & 0 & 2 \\ 2 & 1 & 1 \end{Bmatrix} \quad B = \begin{Bmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \\ 1 & 1 & 3 \end{Bmatrix}$$

$$C = \begin{Bmatrix} 2 & 1 & 3 \\ 2 & 2 & 4 \\ 0 & 1 & 1 \end{Bmatrix}$$

1. Linear transformation of A if vector  $X = (a, b, c)$

solution

$$A = \begin{Bmatrix} 3 & 1 & 2 \\ 1 & 0 & 2 \\ 2 & 1 & 1 \end{Bmatrix}, \quad X = \begin{Bmatrix} a \\ b \\ c \end{Bmatrix}$$

$$T(x) = a \begin{Bmatrix} 3 \\ 1 \\ 2 \end{Bmatrix} + b \begin{Bmatrix} 1 \\ 0 \\ 1 \end{Bmatrix} + c \begin{Bmatrix} 2 \\ 2 \\ 1 \end{Bmatrix}$$

$$T(x) = \begin{Bmatrix} 3a \\ a \\ 2a \end{Bmatrix} + \begin{Bmatrix} b \\ 0 \\ b \end{Bmatrix} + \begin{Bmatrix} 2c \\ 2c \\ c \end{Bmatrix}$$

$$T(x) = \begin{Bmatrix} 3a + b + 2c \\ a + 0 + 2c \\ 2a + b + c \end{Bmatrix}$$

Hence the transformation of

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} \quad \text{gives} \quad \begin{pmatrix} 3a + b + 2c \\ a + 0 + 2c \\ 2a + b + c \end{pmatrix}$$

2. Find the rank of (B+C) transpose

$$B+C = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \\ 1 & 1 & 3 \end{pmatrix} + \begin{pmatrix} 2 & 1 & 3 \\ 2 & 2 & 4 \\ 0 & 1 & 1 \end{pmatrix}$$

$$B+C = \begin{pmatrix} 3 & 1 & 5 \\ 4 & 2 & 5 \\ 1 & 2 & 4 \end{pmatrix}$$

$$(B+C)^T = \begin{pmatrix} 3 & 4 & 1 \\ 1 & 2 & 2 \\ 5 & 5 & 4 \end{pmatrix}$$

To find rank

$$|(B+C)^T| = 3 \begin{vmatrix} 2 & 2 \\ 5 & 4 \end{vmatrix} - 4 \begin{vmatrix} 1 & 2 \\ 5 & 4 \end{vmatrix} + 1 \begin{vmatrix} 1 & 1 \\ 5 & 5 \end{vmatrix}$$

$$= 3(8 - 10) - 4(4 - 10) + 1(5 - 10)$$

$$= 3(-2) - 4(-6) + 1(5 - 10)$$

$$= -6 + 24 - 6$$

$$= 12$$

$$12 \neq 0$$

Hence the Rank of  $(B+C)^T$  is 3.

3. Check whether A, B, and C are singular or non-singular matrix.

For A;

$$|A| = \begin{vmatrix} 3 & 1 & 2 \\ 1 & 0 & 2 \\ 2 & 1 & 1 \end{vmatrix}$$

$$|A| = 3 \begin{vmatrix} 0 & 2 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 2 \\ 2 & 1 \end{vmatrix} + 2 \begin{vmatrix} 1 & 1 \\ 2 & 1 \end{vmatrix}$$

$$\begin{aligned} &= 3(0 - 2) - 1(1 - 4) + 2(1 - 0) \\ &= 3(-2) - 1(-3) + 2(1) \\ &= -6 + 3 + 2 = -1 \\ &-1 \neq 0 \end{aligned}$$

∴ It is a non-singular matrices.

For B;

$$|B| = \begin{vmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \\ 1 & 1 & 3 \end{vmatrix}$$

$$|B| = 1 \begin{vmatrix} 0 & 1 \\ 1 & 3 \end{vmatrix} - 0 \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} + 2 \begin{vmatrix} 2 & 2 \\ 1 & 1 \end{vmatrix}$$

$$\begin{aligned} |B| &= 1(0 - 1) - 0 + 2(2 - 0) \\ &= 1(-1) - 0 + 2(2) \\ &= -1 - 0 + 4 \\ &= 3 \\ &3 \neq 0 \end{aligned}$$

∴ It is a non-singular matrices.

For C;

$$|C| = \begin{vmatrix} 2 & 1 & 3 \\ 2 & 2 & 4 \\ 0 & 1 & 1 \end{vmatrix}$$

$$|C| = 2 \begin{vmatrix} 2 & 4 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 2 & 4 \\ 0 & 1 \end{vmatrix} + 3 \begin{vmatrix} 2 & 2 \\ 0 & 1 \end{vmatrix}$$

$$\begin{aligned} |C| &= 2(2 - 4) - 1(2 - 0) + 3(2 - 0) \\ &= 2(-2) - 1(2) + 3(2) \\ &= -4 - 2 + 6 \\ &= 0 \end{aligned}$$

∴ It is a singular matrix.

18/SCI01/074