

NAME: TENEBE MARLIN-ROSE OMOJI

COURSE: MAT

MATRIC NO: 19/SCI01/086

~~DATE:~~ DEPARTMENT: COMPUTER SCIENCE

$$1. a. M \cdot N = (P\mathbf{i} - 6\mathbf{j} - 8\mathbf{k}) \cdot (4\mathbf{i} + 8\mathbf{j} - \mathbf{k})$$

$$= 4P - 18 + 3$$

$$= 4P - 15$$

For perpendicular vectors

$$= \cancel{4P - 15} \quad 4P - 15 = 0$$

$$4P = 15$$

$$P = 15/4$$

b. M, N and O are coplanar

$$M \cdot (N \times O) = \begin{vmatrix} P & -6 & -3 \\ 4 & 3 & -1 \\ 1 & -3 & 2 \end{vmatrix}$$

$$P \begin{vmatrix} 8 & -1 \\ -3 & 2 \end{vmatrix} + 6 \begin{vmatrix} 4 & -1 \\ 1 & 2 \end{vmatrix} - 3 \begin{vmatrix} 4 & 3 \\ 1 & -3 \end{vmatrix}$$

$$P(6 - 3) + 6(8 + 1) - 3(-12 - 3)$$

$$3P + 54 + 45 = 0$$

$$3P + 99 = 0$$

$$3P = -99$$

$$P = -33$$

$$2. \vec{V} = (3\mathbf{i} + 2\mathbf{j} + 5\mathbf{k}) + (2\mathbf{i} - \mathbf{j} + 6\mathbf{k}) + (5\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})$$

$$\vec{V} = 10\mathbf{i} + 3\mathbf{j} + 8\mathbf{k}$$

$$a_x = 10, \quad a_y = 3, \quad a_z = 8$$

$$|v| = \sqrt{10^2 + 3^2 + 8^2}$$

$$= 13.15$$

The direction cosines are

$$\cos \alpha = \frac{a_x}{|v|} = \frac{10}{13.15} = 0.760$$

$$\cos \beta = \frac{a_y}{|v|} = \frac{3}{13.15} = 0.228$$

$$\cos \gamma = \frac{a_z}{|v|} = \frac{8}{13.15} = 0.608$$

$$\text{Unit vector } \hat{e}_v = \frac{v}{|v|} = \frac{10i + 3j + 8k}{13.15}$$

$$= \frac{10}{13.15}i + \frac{3}{13.15}j + \frac{8}{13.15}k$$

3) If $F = 3u^2i + u^2j + (u+2)k$ and $v = 2u^2i + 3uj + (u-2)k$

$$F \times v = \begin{vmatrix} i & j & k \\ 3u^2 & u^2 & (u+2) \\ 2u & -3u & (u-2) \end{vmatrix}$$

$$i \begin{vmatrix} u^2 & (u+2) \\ -3u & (u-2) \end{vmatrix} - j \begin{vmatrix} 3u^2 & (u+2) \\ 2u & (u-2) \end{vmatrix} + k \begin{vmatrix} -3u & u^2 \\ 2u & -3u \end{vmatrix}$$

$$i(u^3 - 2u + 3u^2 + 6u) - j(3u^2 - 6u - 2u^2 - 4u) + k(-9u^2 - 2u^3)$$

$$i(u^3 + 3u^2 + 4u) - j(u^2 - 10u) + k(-9u^2 - 2u^3)$$

$$\int_0^1 (F \times v) du = \int_0^1 [(u^3 + 3u^2 + 4u)i - (u^2 - 10u)j + (-9u^2 - 2u^3)k] du$$

$$= i \left(\frac{u^4}{4} + \frac{3u^3}{3} + \frac{4u^2}{2} \Big|_0^1 \right) - j \left(\frac{u^3}{3} - \frac{10u^2}{2} \Big|_0^1 \right) + k \left(-\frac{7u^2}{3} - \frac{2u^2}{4} \Big|_0^1 \right)$$

$$= i \left(\frac{u^4}{4} + u^3 + 2u^2 \Big|_0^1 \right) - j \left(\frac{u^3}{3} - 5u^2 \Big|_0^1 \right) + k \left(-\frac{7u^2}{3} - \frac{u^2}{2} \Big|_0^1 \right)$$

$$= i \left[\frac{1}{4} + 1 + 2 - (0) \right] - j \left[\frac{1}{3} - 5 - (0) \right] + k \left[-3 - \frac{1}{2} - (0) \right]$$

$$= i \left[\frac{13}{4} \right] - j \left[\frac{14}{3} \right] + k \left[-\frac{7}{2} \right]$$

$$= \frac{13}{4}i + \frac{14}{3}j - \frac{7}{2}k$$