

NAME: ABDULKADIR ABDULAZIZ AROME

MATRIC NO: 18/MHS01/003

LEVEL: 200 LVL

DEPARTMENT: MEDICINE AND SURGERY

COLLEGE: MEDICINE AND HEALTH SCIENCE

COURSE CODE: ICBS

COURSE TITLE: ANATOMY-EMBRYOLOGY

During the second week of development, the following events occur;

1. •Completion of implantation of the blastocyst
2. •Formation of the bilaminar germ disc(epiblast and hypoblast)
3. •Formation of extraembryonic structures(amniotic
4. cavity, amnion, umbilical vesicle(yolk sac), connecting stalk and chorionic sac)

DAY 8

1. •At the eighth day of development, the blastocyst is partially (slowly) embedded in the endometrium
2. •the syncytiotrophoblast continues to erode the endometrium, thereby eroding endometrial blood vessels and endometrial glands
3. •More cells in the cytotrophoblast divide and migrate into the syncytiotrophoblast, where they fuse and lose their individual cell membranes
4. •The Cells of the inner cell mass or embryoblast also differentiate into 2 layers:
 - a. the hypoblast layer, which is made up of small cuboidal cells, and it is adjacent(nearer) to the blastocyst cavity
 - b. the epiblast layer which is made up of high columnar cells, and it adjacent to the amniotic cavity
5. •The hypoblast and epiblast layers together form a flat ovoid shaped disc called the bilaminar germ disc

6. •At the same time, a small cavity appears within the epiblast which enlarges to form the amniotic cavity
7. •Epiblast cells adjacent to the cytotrophoblast are called amnioblasts
8. •Amnioblasts together with the rest of the epiblast, line the amniotic cavity
9. •The endometrium adjacent to the implantation site is highly vascular

DAY 9

1. •The blastocyst is more deeply embedded in the endometrium, and the penetration defect in the surface epithelium is closed by a coagulum called fibrin
2. •Vacuoles appear at the region of the trophoblast and they fuse to form larger lacunae
3. •this phase of trophoblast development is known as the lacunar stage
4. •the cells of the hypoblast adjacent to the cytotrophoblast form a thin membrane called the exocoelomic (Heuser's) membrane
5. •this membrane lines the inner surface of the cytotrophoblast
6. •the exocoelomic (Heuser's) membrane together with the hypoblast forms the lining of the exocoelomic cavity, or primitive yolk sac or primary umbilical vesicle

DAY 11-12 OF DEVELOPMENT

1. •The blastocyst is **completely** embedded in the endometrium, and the surface epithelium almost entirely covers the original defect in the uterine wall
2. Ruptured capillaries communicate with trophoblastic lacunae, helping us transfer blood, oxygen to developing embryo.
3. •The blastocyst now produces a slight protrusion into the lumen of the uterus
4. •cells of the syncytiotrophoblast penetrate deeper into the stroma and erode the endothelial lining of the endometrial capillaries
5. •These ruptured endometrial capillaries are called sinusoids
6. •The lacunae then begin to communicate with the sinusoids, and maternal blood enters the lacunar system
7. •The communication of the eroded endometrial capillaries with the lacunae establishes the primordial uteroplacental circulation
8. •When maternal blood flows into the lacunae, oxygen and nutritive substances are available to the embryo
9. •a new population of cells appears between the inner surface of the cytotrophoblast and the outer surface of the exocoelomic cavity

10. •These cells which are derived from yolk sac cells form a fine, loose connective tissue called the extraembryonic mesoderm
11. •Soon, large cavities develop in the extraembryonic mesoderm, and when these become confluent, they form a new space known as the extraembryonic cavity or extraembryonic coelom
12. •This space surrounds the primitive yolk sac and amniotic cavity, except where the germ disc is connected to the trophoblast by the connecting stalk (which develops into the umbilical cord)
13. •The extraembryonic mesoderm lining the cytotrophoblast and amnion is called the extraembryonic somatic mesoderm
14. •extraembryonic somatic mesoderm also forms the connecting stalk
15. •the lining covering the yolk sac is known as the extraembryonic splanchnic mesoderm
16. •As the conceptus implants, the endometrial connective tissue cells undergo a transformation, called decidual reaction
17. •During this transformation, the cells of the endometrium swell because of the accumulation of glycogen and lipid in their cytoplasm, and they are known as decidual cells
18. •The primary function of the decidual reaction is to provide nutrition for the early embryo and an immunologically privileged site for the conceptus

DAY 13

1. •The surface defect in the endometrium has been completely covered by the surface epithelium
2. •Occasionally bleeding occurs at the implantation site as a result of increased blood flow into the lacunar spaces
3. •Cells of the cytotrophoblast proliferate locally and penetrate into the syncytiotrophoblast, forming cellular columns surrounded by syncytium
4. •Cellular columns with the syncytial covering are known as primary villi
5. •The primary yolk sac becomes reduced in size and is known as the secondary yolk sac
(This new cavity is known as the secondary yolk sac or definitive yolk sac or the secondary umbilical vesicle)
6. •In humans the yolk sac contains no yolk but is important for the transfer of nutrients between the fetus and mother
7. •This yolk sac is much smaller than the original exocoelomic cavity or primitive yolk sac

8. •During its formation, large portions of the exocoelomic cavity are pinched off to form exocoelomic cysts
9. •Exocoelomic cysts are often found in the extraembryonic cavity or chorionic cavity or extraembryonic coelom
10. •Meanwhile, the extraembryonic coelom expands and forms a large cavity called the chorionic cavity
11. •The extraembryonic mesoderm lining the inside of the cytotrophoblast is then known as the chorionic plate
12. •The only place where extraembryonic mesoderm traverses the chorionic cavity is in the connecting stalk
13. •With development of blood vessels, the connecting stalk becomes the umbilical cord
14. Clinical correlate
15. •The syncytiotrophoblast produces a hormone called the human chorionic gonadotrophin (hCG), which
16. enters the maternal blood via lacunae keeps the corpus luteum secreting estrogens and progesterone □hCG maintains the hormonal activity of the corpus luteum in the ovary during pregnancy
17. •hCG can be detected in maternal blood or urine as early as day 10 of pregnancy and is the basis for
18. pregnancy tests
19. •Enough hCG is produced by the syncytiotrophoblast at the end of the second week to give a positive pregnancy test, even though the woman is probably unaware that she is pregnant
20. •Extrauterine Implantation
21. •Blastocysts may implant outside the uterus
22. •These implantations result in ectopic pregnancies •95% to 98% of ectopic implantations occur in the
23. uterine tubes, most often in the ampulla and isthmus
24. Surface defect in the endometrium has been completely covered by surface epithelium.