

NAME: Gundu-sewuese Oluwashekemi

DEPARTMENT: Nursing

MATRIC NO:19/mhs02/060

Chemistry 102 Assignments

1. CLASSIFICATION OF ALCOHOLS

- I. Classification based on the number of hydrogen atoms attached to the carbon atom containing the hydroxyl group. If the numbers of hydrogen atoms attached to the carbon atom bearing the hydroxyl group are three or two, it is a primary alcohol (1°). If it is one hydrogen atom, it is called a secondary alcohol (2°) and if no hydrogen atom is attached to the carbon atom bearing the hydroxyl group, it is called a tertiary alcohol (3°). Example: $\text{CH}_3\text{CH}_2\text{OH}$ ethanol(1°)
- II. Classification based on the number of hydroxyl groups they possess. Monohydric alcohols have one hydroxyl group present in the alcohol structure. Dihydric alcohol or Glycols have two hydroxyl groups present in the alcohol structure while trihydric alcohols or triols have three hydroxyl groups present in the alcohol structure. Polyhydric alcohols or polyols have more than three hydroxyl groups. Example: $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ propanol (monohydric alcohol).


2. SOLUBILITY OF ALCOHOLS IN WATER AND ORGANIC SOLVENTS

In water, lower alcohols with up to three carbon atoms in their molecules are soluble in water because these lower alcohols can form hydrogen bond with water molecules. The water solubility of alcohols decreases with increasing relative molecular mass.


In organic solvents, all monohydric alcohols are soluble in organic solvents. The solubility of simple alcohols and polyhydric alcohols is largely due to their ability to form hydrogen bonds with water molecules.

3. INDUSTRIAL PREPARATION OF ETHANOL

- i) Carbohydrate such as starch is broken down by diastase contained in malt at a temperature of 60°c to give maltose.




- ii) Maltose is broken down into glucose by maltase found in yeast at a temperature of 15°c to give glucose.



Maltose 15^oc/maltase glucose

iii) Glucose is converted to ethanol at constant temperature of 15^oc by enzyme zymase also contained in yeast.



which contains the enzyme maltase and at a temperature of 15^oc.

$$C_{12}H_{22}O_{11} + H_2O \xrightarrow[15^{\circ}C]{\text{maltase}} 2C_6H_{12}O_6$$

maltose water glucose

STEP 3: The glucose at constant temperature of 15^oc is then converted into alcohol by the enzyme zymase contained also in yeast.

$$C_6H_{12}O_6 \xrightarrow[15^{\circ}C]{\text{zymase}} 2CH_3CH_2OH + 2CO_2 \uparrow$$

Glucose ethanol Carbon(IV)oxide

4 Show the reaction between 2-methylpropanal and butylmagnesium chloride. Hint: Grignard synthesis.

ANSWER

$$CH_3CH(CH_3)CHO + CH_3CH_2CH_2CH_2MgCl \longrightarrow CH_3CH(CH_3)COHCH_2CH_2CH_2CH_3$$

$$\begin{array}{c}
 \text{H} \quad \text{H} \quad \quad \quad \text{O} \\
 | \quad | \quad \quad \quad // \\
 \text{H}-\text{C}-\text{C}-\text{C} \\
 | \quad | \quad \quad \quad \backslash \\
 \text{H} \quad \text{CH}_3 \quad \quad \quad \text{H}
 \end{array}
 +
 \begin{array}{c}
 \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\
 | \quad | \quad | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{MgCl} \\
 | \quad | \quad | \quad | \\
 \text{H} \quad \text{H} \quad \text{H} \quad \text{H}
 \end{array}
 \longrightarrow
 \begin{array}{c}
 \text{H} \quad \text{H} \quad \text{O} \quad \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\
 | \quad | \quad | \quad | \quad | \quad | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\
 | \quad | \quad | \quad | \quad | \quad | \quad | \\
 \text{H} \quad \text{CH}_3 \quad \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \quad \text{H}
 \end{array}$$

2-methyl 3-heptanol

7 Show the reduction reaction of 2-methylpropanal.

ANSWER

i)

$$\begin{array}{c}
 \text{H} \quad \text{C} \\
 | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C} \\
 | \quad | \quad \quad \quad \backslash \\
 \text{H} \quad \text{CH}_3 \quad \quad \quad \text{O}
 \end{array}
 \xrightarrow[\text{CH}_3\text{CH}_2\text{OH}]{\text{NaBH}_4 / \text{LiAlH}_4}
 \begin{array}{c}
 \text{H} \quad \text{H} \quad \text{OH} \\
 | \quad | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\
 | \quad | \quad | \\
 \text{H} \quad \text{CH}_3 \quad \text{H}
 \end{array}$$

2-methyl propanol

OR

ii)

$$\begin{array}{c}
 \text{H} \quad \text{C} \\
 | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C} \\
 | \quad | \quad \quad \quad \backslash \\
 \text{H} \quad \text{CH}_3 \quad \quad \quad \text{O}
 \end{array}
 \xrightarrow[100 \text{ atm}]{\text{H}_2 / \text{Pt} \text{ or Pd or Raney Ni}}
 \begin{array}{c}
 \text{H} \quad \text{H} \quad \text{OH} \\
 | \quad | \quad | \\
 \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\
 | \quad | \quad | \\
 \text{H} \quad \text{CH}_3 \quad \text{H}
 \end{array}$$


4.

8 Propose a scheme for the conversion of propan-1-ol to propan-2-ol.

ANSWER


SCHEME

STEP 1: Dehydration of Propan-1-ol to propene using conc. H_2SO_4




STEP 2: You can use either,

A. Oxymercuration - Demercuration.



Preferable

B. Since propene is asymmetrical, on hydrolysis or addition of water, using a markovnikov procedure, Propan-2-ol can be obtained



You would actually get the 2 products: Propan-1-ol / Propan-2-ol

But following markovnikov's rule, Propan-2-ol would be the major product.