

NAME: Atule Iye Favour

DEPARTMENT: Nursing Science

MATRIC NUMBER: 18/mhs02/049

LEVEL: 200 Level

Pregnancy is a unique period in a woman's lifetime. A number of anatomic, physiologic, biochemical and psychological changes take place. These changes may easily be misinterpreted by physicians who lack experience in regards to pregnancy effects on a woman's body. It is important that physicians caring for women understand the implications of these physiological changes in order to avoid any diagnostic errors and errors of management.

When a woman learns that she is pregnant, her emotions are like roller coaster. To her, she is pregnant. She begins to plan all the things that could be and is in constant motion to await the 9 months until the arrival of her bundle of joy.

SKIN CHANGES

A number of changes take place in the skin of pregnant women. Mechanical stretching of the skin over the abdomen and breasts can lead to striae.



The increased levels of estrogen and progesterone have also been implicated. Usually striae remain permanently with some change in color. Prevention may be achieved with moisturizing creams, especially those containing lanolin and other oily substances. It should be realized, however, that striae may develop despite any preventative measures.

Vascular spider nevi and palmar erythema happen also during pregnancy. There is no clear explanation for these changes, but they most likely represent the result of vasodilatation that happens in the skin during pregnancy. Chloasma and other pigmented lesions can happen as a result of increased melanocyte stimulating hormone activity which in turn is a result of increased estrogen and progesterone levels. These lesions usually begin at about five to six months' gestation. One way that these lesions may be prevented is by the use of screening agents and avoidance of direct sunlight. Skin pruritus affects a number of women and it may be related to increased retention of bile salts in the skin secondary to estrogen effects. Scratching of the skin can then lead to infected excoriations. Local measures with anti-pruritic creams and lotions usually are sufficient.

CHANGES IN THE REPRODUCTIVE SYSTEM

Rhythmic tightening of the uterus occurs as part of preparatory changes for labor. These are called Braxton-Hicks contractions and since the advent of ultrasound, can be seen as early as eight to nine weeks. As the pregnancy advances these contractions become more frequent and they are more likely to be felt by the patient. Usually they happen every 5 to 20 minutes and sometimes they may last as long as 30 minutes.

The genital organs undergo significant changes with increased vascularity of the cervix and increased mucous formation by the cervical glands due to increased levels of estrogen. The vulva and the vagina are also edematous and present increased desquamation and transudation. This leads to an increase in the secretions from the vagina manifesting as increased leucorrhea.

Many patients experience perineal pressure pain, which may be secondary to vascular engorgement of tissues due to estrogen and stasis of blood and to pressure from fetal presenting parts. Pubic pain is also noted and may be secondary to increased joint motility that happens secondary to progesterone's relaxing effect on the pubic symphysis cartilage.

CHANGES IN THE GASTROINTESTINAL SYSTEM

Nausea and vomiting are the most frequent complaints involving the gastrointestinal system and usually happen in early pregnancy while heartburn happen primarily in late pregnancy. The gums become hyperemic and edematous during pregnancy and tend to bleed. The muscular wall of the esophagus is relaxed and this may cause reflux, which in turn can lead to esophagitis and heartburn. The stomach and the intestines have decreased motility presumably due to the effect of progesterone on smooth muscle contractility. This causes an increase in the time that it takes for the stomach to empty. Reduced gastric secretion has also been documented and it could account for the improvement of peptic ulcers sometimes observed in pregnancy.



The liver is affected significantly by pregnancy. Cholestatic jaundice is considered to be the result of estrogen effect on elimination of bilirubin by the liver. The effect of estrogens also, is to increase protein synthesis in the liver, which leads to increased production of fibrinogen and binding proteins. The liver enzymes are usually unaffected with the exception of alkaline phosphatase, which is increased at approximately two fold to four fold that is a result of a placental production. Pregnancy increases the size and decreases the motility of the gall bladder.

CARDIOVASCULAR CHANGES

Of all changes that happen in pregnancy, the single most important is the one involving the cardiovascular system. Adequate cardiovascular adaptation secures good placental development

and thus appropriate fetal growth. The cardiovascular changes involve a substantial change in the blood volume, cardiac output, heart rate, systemic arterial blood pressure, systemic vascular resistance, oxygen consumption and alterations in regional blood flow of various organ systems.

Blood Volume – Significant increases in the blood volume start taking place in the first trimester and continue until the mid-third trimester, at approximately the 32nd to the 34th week. Beyond this point in gestation, the blood volume plateaus. The increase in blood volume with pregnancy appears to serve the essential physiologic needs of both the mother and fetus. It ensures adequate supplies required for normal fetal growth and oxygenation even under circumstances that affect the maternal cardiac output (inferior vena cava compression).

Cardiac Output – It has been well established since the beginning of this century that the cardiac output increases an average of 50 percent during pregnancy. It is generally accepted that cardiac output begins to rise during the first trimester, probably around the tenth week of pregnancy and continues to rise up until the 24th week of gestation. Once it reaches the peak it stays rather stable.

Heart Rate During Normal Pregnancy – The baseline heart rate increases by about 10 to 20 beats per minute. This increase starts early in pregnancy and gradually continues to go upward with the highest values achieved at term. Some investigators, however, suggested that the total increase happens early in pregnancy and remains so throughout the remainder of gestation.


The Heart – A number of changes happen to the heart and are unique to pregnancy. Increasing intra-abdominal contents displace the heart upward with some forward rotation. As a result, the anterior posterior diameter and the cardiothoracic ratio are increased. The overall dimensions of the heart are increased during pregnancy as a result of increased diastolic heart volume without any change in the ventricular wall thickness. Systolic ejection murmurs are common in pregnancy while diastolic murmurs are less frequent. The systolic murmurs are usually the result of the hyper dynamic circulation.

Blood Pressure – A slight decrease in the systolic arterial blood pressure and a significant decrease in the diastolic pressure have been observed to occur in normal pregnancy. This decrease becomes evident in the late first trimester and continues throughout most of the second trimester. The lowest values are noted in mid pregnancy and there after the blood pressure

returns toward non-pregnant levels before term. The degree of change in the blood pressure parameters has been found to be affected by parity, smoking, preexisting hypertension, maternal age and ethnic background. In the typical normal pregnancy, the mean arterial pressure (diastolic plus 1/3 of the difference between systolic and diastolic) is less than 85 mm of mercury. Studies have found that when the mean arterial blood pressure in the mid second trimester is higher than 90 mm of mercury, there is increased perinatal mortality and morbidity.

BREAST CHANGES

- Increased size of the breasts
- Marked hypertrophy and proliferation of the ducts (oestrogen and progesterone)
- Vascularity is increased
- The nipples become larger, erectile and deeply pigmented



Finally, a number of women may experience backache in the upper back, which is secondary to muscle tension from increasing breast size and discomfort. Most women, however, experience low back pain secondary to muscular fatigue and strain that is caused by the changes in body balance from the growing uterus. Several patients also may experience pressure on nerve roots that in turn may lead to muscular spasms and pelvic joint pains secondary to bone ligament relaxation from the sex hormones. The changes that happen on the ligaments and the cartilage of the pelvic bones secondary to the sex hormones may also lead some women to

present with gait alterations. Finally, a number of women may experience paresthesia (numbness and tingling of fingers and toes). A number of theories are suggested for the explanation of these symptoms. The fingers and upper extremities are effected if lordosis posture is extreme; the head and neck are flexed, putting strain on the brachial nerves and causing tingling of hands and arms. Toes and lower extremities are affected if gravid uterus presses on femoral veins and nerves supplying lower extremities, thus interfering with circulation and causing paresthesia. Edema may cause pressure and tingling of hands or feet, especially in hands when rising in the morning. Sometimes excessive edema of the hands may lead to carpal tunnel syndrome. Finally, Vitamin B deficiency, hypoglycemia and hyperventilation have been suggested as causes of these symptoms.