

NAME: EDIGBE JOY ELOHOR

MATRIC NUMBER: 18/MHS02/065

DEPARTMENT: NURSING

COURSE CODE: PHS 212

**Assignment:- Discuss the factors affiliating the movement of sperm in the female reproductive tract.**

At coitus, human sperm are deposited into the anterior vagina, where, to avoid vaginal acid and immune responses, they quickly contact cervical mucus and enter the cervix. Cervical mucus filters out sperm with poor morphology and motility and as such only a minority of ejaculated sperm actually enter the cervix. In the uterus, muscular contractions may enhance passage of sperm through the uterine cavity. A few thousand sperm swim through the uterotubal junctions to reach the Fallopian tubes (uterine tubes, oviducts) where sperm are stored in a reservoir, or at least maintained in a fertile state, by interacting with endosalpingeal (oviductal) epithelium. As the time of ovulation approaches, sperm become capacitated and hyper activated, which enables them to proceed towards the tubal ampulla. Sperm may be guided to the oocyte by a combination of thermotaxis and chemotaxis.

Motility hyper activation assists sperm in penetrating mucus in the tubes and the cumulus oophorus and zona pellucida of the oocyte, so that they may finally fuse with the oocyte plasma membrane. Knowledge of the biology of sperm transport can inspire improvements in artificial insemination, IVF, the diagnosis of infertility and the development of contraceptives. Passage of sperm through the female reproductive tract is regulated to maximize the chance of fertilization and ensure that sperm with normal morphology and vigorous motility will be the ones to succeed.

Oocytes are usually fertilized within hours of ovulation. In humans, there is evidence that fertilization occurs when intercourse takes place up to five days before ovulation. Because sperm are terminally differentiated cells, deprived of an active transcription and translation apparatus, they must survive in the female without benefit of reparative mechanisms available to many other cells. Sperm are subjected to physical stresses during ejaculation and contractions of the female tract, and they may sustain oxidative damage.

Furthermore, because sperm are allogenic to the female, they may encounter the defenses of the female immune system meant for infectious organisms. Thus, sperm must somehow use their limited resources to maintain their fertility in the face of numerous impediments. As it is, of the millions of sperm inseminated at coitus in humans, only a few thousand reach the Fallopian tubes and, ordinarily, only a single sperm fertilizes an oocyte.