

NAME: OKEREKE OGETCHI ANTHONIA
 DEPARTMENT: PHARMACY
 COURSE CODE: MAT104
 MATRIC NO: 191MHS1103

COVID-19 ASSIGNMENT

1. $y = \frac{1}{x-2}$ state the domain and codomain
 \rightarrow The function is defined for all real numbers except 2.
 \rightarrow The co-domain is the set of real numbers except $y=0$.
 \rightarrow The domain is the set of real numbers except 2.

2. $K = \ln v$
 $\frac{dk}{dv} = \frac{1}{v}$
 3a. $2x - 3y - 2 = 0$
 $-3y = 2 - 2x$
 $y = \frac{2-2x}{-3}$
 $y = \frac{2x-2}{3}; \frac{2(x-1)}{3}$

b. $x^2 + y^2 = 4$
 $y^2 = 4 - x^2$
 $y = \pm \sqrt{4-x^2}$

4. $P = \sin^{-1} t$, find the derivative of P.
 $P = t$ $t = \sin P$ $\frac{d}{dt}$
 \sin
 $\frac{dt}{dp} = \cos P; \frac{dp}{dt} = \frac{1}{\cos P}$
 Recall, $\cos^2 y + \sin^2 y = 1$
 $\cos y = \pm \sqrt{1 - \sin^2 y}$
 But $t = \sin P$, $\cos P = \sqrt{1-t^2}$
 $\therefore \frac{dP}{dt} = \frac{1}{\sqrt{1-t^2}}$

5. If $f(x) = 2x^2 - 5$ and $g(x) = 4x - 2$, find $f \circ g(x)$ and $g \circ f(x)$
 $f(x) = 2x^2 - 5$; $g(x) = 4x - 2$
 $f \circ g(x) = 2(4x-2)^2 - 5$
 $= 2(16x^2 - 16x + 4) - 5$
 $= 32x^2 - 32x + 8 - 5$
 $= 32x^2 - 32x + 3$
 $g \circ f(x) = 4(2x^2 - 5) - 2$
 $= 8x^2 - 20 - 2$
 $= 8x^2 - 22$

Q. Show that $f(x) = f_0(x) + f_1(x)$

If $f(x) = 3x^2 - 2x + 1 = 0$

$f_0(x) = f(x) + f(-x)$

$f(-x) = 3(-x)^2 - 2(-x) + 1$
 $= 3x^2 + 2x + 1$

$f_0(x) = \frac{3x^2 - 2x + 1 + 3x^2 + 2x + 1}{2}$

$= \frac{6x^2 + 2}{2} = 3x^2 + 1$

$f_1(x) = \frac{3x^2 - 2x + 1 - (3x^2 + 2x + 1)}{2}$

$= \frac{-4x}{2} = -2x$

$f_0(x) + f_1(x) = 3x^2 + 1 - 2x$
 $= 3x^2 - 2x + 1$

Q. Differentiate $y = \cos x$ according to 1st principle

$y = \cos x$

$y + \delta y = \cos(x + \delta x)$

Subtract y from both sides

$\delta y = \cos(x + \delta x) - y$

but $y = \cos x$

$\therefore \delta y = \cos(x + \delta x) - \cos x \dots \dots \text{---} \text{---}$

consider from trig

$\cos(A - B)$

$\cos(A + B) = \cos A \cos B - \sin A \sin B$

$\cos(A - B) = \cos A \cos B + \sin A \sin B$

$\cos(A + B) - \cos(A - B) = -2 \sin A \sin B$

Compare (1) and (2)

$A + B = x + \delta x \dots \dots \text{---}$

$A - B = x \dots \dots \text{---}$

Adding equation (1) & (2)

$2A = 2x + \delta x$

$A = \frac{2x + \delta x}{2} = \frac{2x}{2} + \frac{\delta x}{2}$

$A = \frac{2x + \delta x}{2} \quad A = \frac{2x}{2} + \frac{\delta x}{2}$

If $A = x + \frac{\delta x}{2}$

From eq (1)

$B = A - x$

$B = \frac{x + \frac{\delta x}{2}}{2} - x$

$B = \frac{\frac{\delta x}{2} + x - 2x}{2}$

$B = \frac{\delta x}{2}$

compare eq (1) and (2)

$\cos(x + \frac{\delta x}{2}) - \cos x = -2 \sin(\frac{x + \frac{\delta x}{2}}{2})$

$\sin(\frac{dx}{2})$

$$\frac{dy}{dx} = 6t \div \frac{-2}{t^3}$$

$$= 6t \times \frac{t^3}{-2}$$

$$= 3t \times t^3$$

$$= 3t^4$$

$$= -3t^4$$

$$= -3t^4$$

9. Find $\frac{dy}{dx}$ if $y = x^2 \cos 2x e^{4x}$

Taking log of both sides

$$\ln y = \ln x^2 + \ln \cos 2x + \ln e^{4x}$$

Differentiating w.r.t x

$$\frac{1}{y} \frac{dy}{dx} = \frac{2}{x} + 1(-2 \sin 2x) + 4 \cos 2x$$

$$\frac{1}{y} \frac{dy}{dx} = \frac{2}{x} - 2 \sin 2x + 4 \cos 2x$$

Multiplying both sides by y

$$\frac{dy}{dx} = y \left(\frac{2}{x} - 2 \sin 2x + 4 \right)$$

$$\text{But } y = x^2 \cos 2x e^{4x}$$

$$\frac{dy}{dx} = x^2 \cos 2x e^{4x} \times \left(\frac{2}{x} - 2 \sin 2x + 4 \right)$$

$$\frac{dy}{dx} = -2 \sin \left(x + \frac{8x}{2} \right) \sin \left(\frac{8x}{2} \right)$$

$$\frac{dy}{dx} = -2 \sin \left(x + \frac{8x}{2} \right) \sin \left(\frac{8x}{2} \right)$$

$$= -\sin \left(x + \frac{8x}{2} \right) \sin \left(\frac{8x}{2} \right)$$

$$\frac{dy}{dx} = -\sin \left(x + \frac{8x}{2} \right) \sin \left(\frac{8x}{2} \right) \dots \dots \dots$$

A standard limit

$$\lim_{8x \rightarrow 0} \frac{\sin \left(\frac{8x}{2} \right)}{\frac{8x}{2}} = 1$$

$$8x \rightarrow 0$$

Find limit (4th) as $8x \rightarrow 0$

$$\lim_{dx} \frac{dy}{dx} = \lim_{dx} -\sin \left(x + \frac{8x}{2} \right) \frac{\sin \left(\frac{8x}{2} \right)}{\frac{8x}{2}}$$

$$= -\sin(x+0) \cdot 1$$

$$= -\sin(x)$$

$$\lim_{8x \rightarrow 0} \frac{dy}{dx} = -\sin(x)$$

$$8x \rightarrow 0$$

8. If $y = 3t^2$ and $x = \frac{1}{t^2}$ Find $\frac{dy}{dx}$

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$$

$$= \frac{dy}{dt} \div \frac{dx}{dt}$$

$$\frac{dy}{dt} = 6t \quad ; \quad \frac{dx}{dt} = -\frac{2}{t^3}$$

10. Given that $y = \sin(3x^3 + 5)$ find the derivative of y

$$y = \sin(3x^3 + 5)$$

$$\text{let } u = 3x^3 + 5 : \frac{du}{dx} = 9x^2$$

$$y = \sin u : \frac{dy}{du} = \cos u$$

$$\therefore \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$
$$= \cos u \times 9x^2$$

$$\frac{dy}{dx} = 9x^2 \cos u$$

$$\text{but } u = 3x^3 + 5$$

$$\frac{dy}{dx} = 9x^2 \cos(3x^3 + 5)$$