

**NAME: ENECHUKWU CHIDUMEBI.W**

**MATRIC NO: 18/MHS02/074**

**DEPARTMENT: NURSING**

**ELUCIDATE THE PHYSIOLOGICAL ADAPTATIONS OF THE FEMALE TO PREGNANCY**

Maternal physiological changes in pregnancy are the adaptation during pregnancy that a woman's body undergoes to accommodate the growing embryo or fetus. The physiologic changes are entirely normal, and include behavioral (brain), cardiovascular (heart and blood vessel), hematologic (blood), metabolic, renal (kidney), posture, and respiratory (breathing) changes. Increases in blood sugar, breathing, and cardiac output are all expected changes that allow a pregnant woman's body to facilitate the proper growth and development of the embryo or fetus during the pregnancy.

The pregnant woman and the placenta also produce many other hormones that have a broad range of effects during the pregnancy.

**Hormonal**

Pregnant women experience numerous adjustments in their endocrine system that help support the fetus. The fetal-placental unit secretes steroid hormones and proteins that alter the function of various maternal endocrine glands. Sometimes, the changes in certain hormone levels and their effects on their target organs can lead to gestational diabetes and gestational hypertension.

**Fetal- placental unit**

Levels of progesterone and estrogen rise continually throughout pregnancy, suppressing the hypothalamic axis and subsequently the menstrual cycle. Progesterone is the first produced by the corpus luteum and then by the placenta in the second trimester. Women also experience increased human chorionic gonadotropin ( $\beta$ -hCG), which is produced by the placenta.

**Pancreatic insulin**

The placenta also produces human placental lactogen (hPL), which stimulates maternal lipolysis and fatty acid metabolism. As a result, this converts blood glucose for use by the fetus. It can also decrease maternal tissue sensitivity to insulin, resulting in gestational diabetes.

**Pituitary gland**

The pituitary gland grows by about one-third as a result of hyperplasia of the lactotrophs in response to the high plasma estrogen. Prolactin, which is produced by the lactotrophs, increases progressively throughout pregnancy. Prolactin mediates a change in the

structure of the breast mammary glands from ductal to lobular-alveolar and stimulates milk production.

### **Parathyroid**

Fetal skeletal formation and then later lactation challenges the maternal body to maintain their calcium levels. The fetal skeleton requires approximately 30 grams of calcium by the end of pregnancy. The mother's body adapts by increasing parathyroid hormone, leading to an increased calcium reabsorption by the kidneys. Maternal total serum calcium decreases due to maternal hypoalbuminemia, but the ionized calcium levels are maintained.

### **Adrenal glands**

Total cortisol increases to three times of non-pregnant levels by the third trimester. The increased estrogen in pregnancy leads to increased corticosteroid-binding globulin production and in response the adrenal gland produces more cortisol. The net effect is an increase of free cortisol. This contributes to insulin resistance of pregnancy and possibly striae. Despite the increase in cortisol, the pregnant mom does not exhibit crushing syndrome or symptoms of high cortisol. One theory is that high progesterone levels acts as an antagonist to the cortisol.

The adrenal gland also produces more aldosterone, leading to an eight-fold increase in aldosterone. Women do not show signs of hyperaldosterone, such as hypokalemia, hypernatremia, or high blood pressure.

The adrenal gland also produces more androgens, such as testosterone, but this is buffered by estrogen's increase in sex-hormone binding globulin (SHBG). SHBG binds avidly to testosterone and a lesser degree DHEA.

### **Thyroid**

The thyroid enlarges and may be more easily felt during the first trimester. The increase in kidney clearance during pregnancy causes more iodide to be excreted and causes relative iodine deficiency and as a result an increase in thyroid-binding globulin (TBG) leads to an increase in total thyroxine (T4), but free thyroxine (T4) and triiodothyronine (T3) remain normal.

### **Breast size**

A woman's breasts grow during pregnancy, usually 1 to 2 cup sizes and potentially several cup sizes. A woman who wore a cup bra prior to her pregnancy may need to buy an F cup or larger bra while nursing. A woman's torso may increase one or two sizes. An average of 80% of women wear the wrong bra size, and mothers who are preparing to nurse can benefit from a professional bra fitting from a lactation consultant. Once the baby is born up to about 50–73

hours after birth, the mother will experience her breasts filling with milk (sometimes referred to as “the milk coming in. Once lactation begins, the woman's breasts swell significantly and can feel achy, lumpy and heavy (which is referred to as engorgement). Her breasts may increase in size again by an additional 1 or 2 cup sizes, but individual breast size may vary depending on how much the infant nurses from each breast.

## **Cardiovascular**

The heart adapts to the increased cardiac demand that occurs during pregnancy in many ways.

- Cardiac output (Lit./Min.): 6.26
- Stroke Volume (Ml.): 75
- Heart Rate (Per min.): 85
- Blood Pressure: Unaffected

Cardiac output increases throughout early pregnancy, and peaks in the third trimester, usually to 30-50% above baseline. Estrogen mediates this rise in cardiac output by increasing the pre-load and stroke volume, mainly via a higher overall blood volume (which increases by 40–50%). The heart rate increases, but generally not above 100 beats/ minute. Total systematic vascular resistance decreases by 20% secondary to the vasodilatory effect of progesterone. Overall, the systolic and diastolic blood pressure drops 10–15 mm Hg in the first trimester and then returns to baseline in the second half of pregnancy. All of these cardiovascular adaptations can lead to common complaints, such as palpitations, decreased exercise tolerance, and dizziness.

## **Hematology**

### **Blood volume and hemoglobin concentration**

During pregnancy the plasma volume increases by 40-50% and the red blood cell volume increases only by 20–30%. These changes occur mostly in the second trimester and prior to 32 weeks gestation. Due to dilution, the net result is a decrease in hematocrit or hemoglobin, which are measures of red blood cell concentration. Erythropoietin, which stimulates red blood cell production, increases throughout pregnancy and reaches approximately 150 percent of their pregnancy levels at term. The slight drop in hematocrit or hemoglobin is most pronounced at the end of the second trimester and slowly improves when reaching term.

### **Platelet and white cell count**

The effect of pregnancy on platelet count is unclear, with some studies demonstrating a mild decline in platelet count and other studies that show no effect. The white blood cell count increases with occasional appearance of myelocytes or metamyelocytes in the blood. During labor, there is a rise in leukocyte count.

### **Hypercoagulability**

A pregnant woman will also become hypercoagulable, leading to increased risk for developing blood clots and embolisms, such as deep vein thrombosis and pulmonary embolism. Women

are 4-5 times more likely to develop a clot during pregnancy and in the postpartum period than when they are not pregnant. Hypercoagulability in pregnancy likely evolved to protect women from hemorrhage at the time of miscarriage or childbirth. In third world countries, the leading cause of maternal death is still hemorrhage. In the United States 2011-2013, hemorrhage made up of 11.4% and pulmonary embolisms made up of 9.2% of all pregnancy-related deaths.

The increased risk of clots can be attributed to several things. Plasma levels of pro-coagulation factors increased markedly in pregnancy, including: von Willebrand Factor, fibrinogen, factor VII, factor VIII, and factor X. Both the production of prostacyclin (an inhibitor of platelet aggregation) and thromboxane (an inducer of platelet aggregation and a vasoconstrictor) are increased, but overall there is an increase in platelet reactivity which can lead to a predisposition to clots. There is also increased blood stasis due to the compression of the vena cava by the enlarging uterus. Many factors have been shown to increase the risk of clots in pregnancy, including baseline thrombophilia, cesarean section, preeclampsia, etc. Clots usually develop in the left leg or the left iliac/ femoral venous system. Recently, there have been several case reports of May-Thurner Syndrome in pregnancy, where the right common iliac artery compresses the below left common iliac vein.

### **Edema**

Edema, or swelling, of the feet is common during pregnancy, partly because the enlarging uterus compresses veins and lymphatic drainage from the leg.

---

### **Metabolic**

During pregnancy, both protein metabolism and carbohydrate metabolism are affected. One kilogram of extra protein is deposited, with half going to the fetus and placenta, and another half going to uterine contractile proteins, breast glandular tissue, plasma protein, and haemoglobin.

An increased requirement for nutrients is given by fetal growth and fat deposition. Changes are caused by steroid hormones, lactogen, and cortisol.

Maternal insulin resistance can lead to gestational diabetes. Increased liver metabolism is also seen, with increased gluconeogenesis to increase maternal glucose levels.

### **Body weight**

Some degree of weight gain is expected during pregnancy. The enlarging uterus, growing fetus, placenta, amniotic fluid, normal increase in body fat, and increase in water retention all contribute weight gain during pregnancy. The amount of weight gain can vary from 5 pounds (2.3 kg) to over 100 pounds (45 kg). In the United States, the range of weight gain that doctors generally recommend is 25 pounds (11 kg) to 35 pounds (16 kg), less if the woman is overweight, more (up to 40 pounds (18 kg)) if the woman is underweight.

### **Nutrition**

Nutritionally, pregnant women require a caloric increase of 350 kcal/day and an increase in protein to 70 or 75 g/day. There is also an increased folate requirement from 0.4 to 0.8 mg/day (important in preventing neural tube defects). On average, a weight gain of 20 to 30 lb (9.1 to 13.6 kg) is experienced.

All patients are advised to take prenatal vitamins to compensate for the increased nutritional requirements. The use of Omega 3 fatty acids supports mental and visual development of infants. Choline supplementation of research mammals supports mental development that lasts throughout life.

### **Renal and lower reproductive tract**

Progesterone causes many changes to the genitourinary system. A pregnant woman may experience an increase in the size of the kidneys and ureter due to the increase blood volume and vasculature. Later in pregnancy, the woman might develop physiological hydronephrosis and hydroureter, which are normal. Progesterone causes vasodilatation and increased blood flow to the kidneys, and as a result glomerular filtration rate (GFR) commonly increases by 50%, returning to normal around 20 weeks postpartum. The increased GFR increases the excretion of protein, albumin, and glucose. The increased GFR leads to increased urinary output, which the woman may experience as increased urinary frequency. Progesterone also causes decreased motility of the ureters, which can lead to stasis of the urine and hence an increased risk of urinary tract infection. Pregnancy alters the vaginal microbiota with a reduction in species/genus diversity. Physiological hydronephrosis may appear from six weeks.

### **Gastrointestinal**

Changes in the gastrointestinal (GI) system during pregnancy are caused by the enlarging uterus and hormonal changes of pregnancy. Anatomically, the intestine and stomach are pushed up from their original positions by the enlarging uterus. While there aren't any intrinsic changes in the sizes of the GI organs, the portal vein increases in size due to the hyperdynamic state of pregnancy. Elevated levels of progesterone and estrogen mediate most of the functional changes of the GI system during pregnancy. Progesterone causes smooth muscle relaxation which slows down GI motility and decreases lower esophageal sphincter (LES) tone. The resulting increase in intragastric pressure combined with lower LES tone leads to the gastroesophageal reflux commonly experienced during pregnancy.

The increased occurrence of gallstones during pregnancy is due to inhibition of gallbladder contraction (as result of increased smooth muscle relaxation mediated by progesterone) and reduced biliary transportation of bile (mediated by estrogen) which results in cholestasis of pregnancy.

Nausea and vomiting of pregnancy, commonly known as “morning sickness”, is one of the most common GI symptoms of pregnancy. It begins between the 4 and 8 weeks of pregnancy and usually subsides by 14 to 16 weeks. The exact cause of nausea is not fully understood but it correlates with the rise in the levels of human chorionic gonadotropin, progesterone, and the resulting relaxation of smooth muscle of the stomach. Hyperemesis gravidarum, which is a severe form of nausea and vomiting of pregnancy can lead to nutritional deficiencies, weight

loss, electrolytes imbalance and is one of the leading causes of hospitalization in the first trimester of pregnancy.

Constipation is another GI symptom that is commonly encountered during pregnancy. It is associated with the narrowing of the colon as it gets pushed by the growing uterus found adjacent it leading to mechanical blockade. Reduced motility in the entire GI system as well as increased absorption of water during pregnancy are thought to be contributing factors.

Dietary cravings and dietary as well as olfactory avoidance of certain types of food are common in pregnancy. Although the exact mechanisms of these symptoms are not fully explained, it is thought that dietary cravings may arise from the thought that certain foods might help relieve nausea. Pica, which is the intense craving for unusual materials such as clay and ice has also been reported in pregnancy.

Hemorrhoids and gingival disease are two common pregnancy associated physical findings involving the gastrointestinal system. Hemorrhoids arise as a result of constipation and venous congestion that are common in pregnancy. Gingival disease is thought to be related to gum softening and edema (swelling from fluid collection) that is mostly observed in pregnancy. The mechanism and reason for the gingival changes are poorly understood.

### **Immune intolerance**

The fetus inside a pregnant woman may be viewed as an unusually successful allograft, since it genetically differs from the woman. In the same way, many cases of spontaneous abortion may be described in the same way as maternal transplant rejection.

### **Musculoskeletal**

Neuromechanical adaptations to pregnancy refers to the change in gait, postural parameters, as well as sensory feedback, due to the numerous anatomical, physiological, and hormonal changes women experience during pregnancy. Such changes increase their risk for musculoskeletal disorders and fall injuries. Musculoskeletal disorders include lower-back pain, leg cramps, and hip pain. Pregnant women fall at a similar rate (27%) to women over age of 70 years (28%). Most of the falls (64%) occur during the second trimester. Additionally, two-thirds of falls are associated with walking on slippery floors, rushing, or carrying an object. The root causes for these falls are not well known. However, some factors that may contribute to these injuries include deviations from normal posture, balance, and gait.

The body's posture changes as the pregnancy progresses. The pelvis tilts and the back arches to help keep balance. Poor posture occurs naturally from the stretching of the woman's abdominal muscles as the fetus grows. These muscles are less able to contract and keep the lower back in proper alignment. The pregnant woman has a different pattern of gait. The step lengthens as the pregnancy progresses, due to weight gain and changes in posture. On average, a woman's foot can grow by a half size or more during pregnancy. In addition, the increased body weight of pregnancy, fluid retention, and weight gain lowers the arches of the foot, further adding to the foot's length and width. The influences of increased hormones such as estrogen and relaxin initiate the remodeling of soft tissues, cartilage and ligaments. Certain skeletal joints such as the pubic symphysis and sacroiliac widen or have increased laxity

The addition of mass, particularly around the torso, naturally changes a pregnant mother's center of mass (COM). The change in COM requires pregnant mothers to adjust their bodies to maintain balance.