

Okunnu Ifedola Rachel

18/MHS/07/039

Pharmacology

BCH204 VITAMINS AND COENZYME

1a) Coenzymes are small molecules. They cannot by themselves catalyze a reaction but they can help enzymes to do so. In technical terms, coenzymes are organic nonprotein molecules that bind with the protein molecule (apoenzyme) to form the active enzyme (holoenzyme).

1b)

Fat soluble vitamins	Water soluble vitamins
----------------------	------------------------

Soluble In fat	Soluble in water
----------------	------------------

Cannot work as co-enzymes	Work as co-enzymes
---------------------------	--------------------

Carrier proteins are present	No carrier protein required
------------------------------	-----------------------------

Stored majorly in the liver	Has no storage
-----------------------------	----------------

Absorption occurs along with lipids and require me bile salt	Absorption simple
--	-------------------

1c)

Niacin is a coenzyme, like thiamine and riboflavin, that is responsible for energy release from carbohydrates. A niacin deficiency can lead to pellagra, a disabling disease with symptoms that may be characterized by four “Ds”: depression, diarrhea, delirium and dementia.

Niacin is found in fortified breads and cereals. Protein foods, such as eggs, fish, meat, dairy milk and poultry, are naturally rich in niacin. They are also plentiful in the amino acid tryptophan, which can be synthesized into niacin by the liver. Chicken breast, ground beef, halibut, tuna and turkey are particularly good sources of tryptophan. In the vegetable kingdom, asparagus, baked potatoes and cantaloupe have significant amounts of tryptophan.

Niacin has been used to lower LDL cholesterol and raise HDL cholesterol when administered as a drug under medical guidance. In heavy doses, niacin has been known to cause a “niacin flush” due to the capillaries increasing in size. This condition can lead to fatigue and even liver damage. Caution should be used if one is taking niacin or B-complex supplements.