

Jeremiah Judith Chidiebere

18/mhs02/098

**Course: human
physiology**

Course code: Phs 212

**Question: discuss the
factors facilitating the
movement of sperm in
the female reproductive
tract**

At coitus, human sperm are deposited into the anterior vagina, where, to avoid vaginal acid and immune responses, they quickly contact cervical mucus and enter the cervix.

Cervical mucus filters out sperm with poor morphology and motility and as such only a minority of ejaculated sperm actually enter the cervix. In the uterus, muscular contractions may enhance passage of sperm through the uterine cavity. A

few thousand sperm swim through the uterotubal junctions to reach the Fallopian tubes (uterine tubes, oviducts) where sperm are stored in a reservoir, or at least maintained in a fertile state, by interacting with endosalpingeal (oviductal) epithelium. As the time of ovulation approaches, sperm become capacitated and hyperactivated, which enables them to proceed towards the tubal ampulla. Sperm may be guided to the oocyte by a combination of thermotaxis and chemotaxis. Motility hyperactivation assists sperm in penetrating mucus in the tubes and the cumulus oophorus and zona pellucida of the oocyte, so that they may finally fuse with the oocyte plasma membrane. Knowledge of the biology of sperm transport can inspire

improvements in artificial
insemination, IVF, the diagnosis
of infertility and the
development of contraceptives.