

Onwughalu Chiamaka Vivian

18/MHS07/044

PHS 212

Assignment: Discuss the physiology of balance

Answer:

The Physiology Of Balance: Vestibular Function

- balance is mediated by the vestibular nuclei in the brain stem
- the labyrinth (a part of the inner ear), is a major organ of our vestibular (balance) system
 - the three semicircular canals of the labyrinth is associated with sensing rotary motion (1)
 - the brain senses the direction and speed of rotation of the head by the movement of fluid in the semicircular canals (1)
 - balance is maintained by the interactions between the labyrinth and other systems in the body, such as the visual and skeletal systems (1)
 - the main inputs into the balance system are the:
 - vestibular labyrinths
 - visual system (eyes)
 - somatosensory system, especially proprioception
 - the main outputs from the vestibular nuclei are:
 - vestibulo-ocular:
 - permitting reflex eye movements related to posture
 - vestibulo-spinal which supply:
 - anti-gravity muscles in the lower limbs
 - reflex arcs which control gait

The vestibular system is the sensory apparatus of the inner ear that helps the body maintain its postural equilibrium. The information furnished by the vestibular system is also essential for coordinating the position of the head and the movement of the eyes. There are two sets of end organs in the inner ear, or labyrinth: the semicircular canals, which respond to rotational movements (angular acceleration); and the utricle and saccule within the vestibule, which respond to changes in the position of the head with respect to gravity (linear acceleration). The information these organs deliver is proprioceptive in character, dealing with events within the body itself, rather than exteroceptive, dealing with events outside the body, as in the case of the responses of the cochlea to sound. Functionally these organs are closely related to the cerebellum and to the reflex centres of the spinal cord and brainstem that govern the movements of the eyes, neck, and limbs.

In vertebrates the utricular maculae in the inner ear contain an otolithic membrane and otoconia (particles of calcium carbonate) that bend hair cells in the direction of gravity. This response to gravitational pull helps animals maintain their sense of balance.

Although the vestibular organs and the cochlea are derived embryologically from the same formation, the otic vesicle, their association in the inner ear seems to be a

matter more of convenience than of necessity. From both the developmental and the structural point of view, the kinship of the vestibular organs with the lateral line system of the fish is readily apparent. The lateral line system is made up of a series of small sense organs located in the skin of the head and along the sides of the body of fishes. Each organ contains a crista, sensory hair cells, and a cupula, as found in the ampullae of the semicircular ducts. The cristae respond to waterborne vibrations and to pressure changes.

Disturbances of the vestibular system

The relation between the vestibular apparatus of the two ears is reciprocal. When the head is turned to the left, the discharge from the left horizontal canal is decreased, and vice versa. Normal posture is the result of their acting in cooperation and in opposition. When the vestibular system of one ear is damaged, the unrestrained activity of the other causes a continuous false sense of turning (vertigo) and rhythmical, jerky movements of the eyes (nystagmus), both toward the uninjured side. When the vestibular hair cells of both inner ears are injured or destroyed, as can occur during treatment with the antibiotics gentamicin or streptomycin, there may be a serious disturbance of posture and gait (ataxia) as well as severe vertigo and disorientation. In younger persons the disturbance tends to subside as reliance is placed on vision and on proprioceptive impulses from the muscles and joints as well as on cutaneous impulses from the soles of the feet to compensate for the loss of information from the semicircular canals. Recovery of some injured hair cells may occur.

Routine tests of vestibular function traditionally have involved stimulation of the semicircular canals to elicit nystagmus and other vestibular ocular reflexes. Rotation, which can cause vertigo and nystagmus, as well as temporary disorientation and a tendency to fall, stimulates the vestibular apparatus of both ears simultaneously. Because otoneurologists are usually more interested in examining the right and left ears separately, they usually employ temperature change as a stimulant. Syringing the ear canal with warm water at 44 °C (111 °F) or with cool water at 30 °C (86 °F) elicits nystagmus by setting up convection currents in the horizontal canal. The duration of the nystagmus may be timed with a stopwatch, or the rate and amplitude of the movements of the eyes can be accurately recorded by picking up the resulting rhythmical variations in the corneoretinal direct current potentials, using electrodes pasted to the skin of the temples—a diagnostic process called electronystagmography. An abnormal vestibular apparatus usually yields a reduced response or no response at all.

The vestibular system may react to unaccustomed stimulation from the motion of an aircraft, a ship, or a land vehicle to produce a sense of unsteadiness, abdominal discomfort, nausea, and vomiting. Effects not unlike motion sickness, with vertigo and nystagmus, can be observed in the later stages of acute alcoholic intoxication. Vertigo accompanied by hearing loss is a prominent feature of the periodic attacks experienced by patients with Ménière disease, which, until the late 19th century, was confused with epilepsy. It was referred to as apoplectiform cerebral congestion and

was treated by purging and bleeding. Other forms of vertigo may present the otoneurologist with more difficult diagnostic problems.