

Name: Opoola Temitope Susan

Matric No: 18/MHS02/164

Department: Nursing

Course code: PHS212


Date: Tuesday 19th, June, 2020

Elucidate the pathway involved in Taste

The tongue is a muscular organ in the mouth. The tongue is covered with moist, pink tissue called mucosa. Tiny bumps called papillae give the tongue its rough texture. Thousands of taste buds cover the surfaces of the papillae. Taste buds are collections of nerve-like cells that connect to nerves running into the brain.

The tongue contains small bumps called papillae, within or near which taste buds are situated. In the tongue's taste buds, the taste receptors receive sensory input via two important mechanisms – depolarization and neurotransmitter release. Intake of salty foods leads more sodium ions to enter the receptor, causing the said mechanisms. The same is true with intake of sour foods (hydrogen ions) and sweet foods (sugar molecules), both of which result to the closing of K^+ channels upon their entry.

From the axons of the taste receptors, the sensory information is transferred to the three taste pathways via the branches of cranial nerves VII, IX and X. The chorda tympani of CN VII (facial nerve) carries the taste sensory input from the tongue's anterior two-thirds. Then, the rest of the taste sensations from the throat, palate and posterior tongue are transmitted by the branches of CN IX (glossopharyngeal nerve) and CN X (vagus nerve). From these cranial nerves, taste sensory input travels through the nerve fiber synapses to the solitary tract, the ventral posteromedial thalamic nuclei, and the thalamus. In these three locations, there are clustered neurons which respond to the same taste (sweet, sour, salty or bitter). The thalamus relays the information to the primary gustatory cortex located in the somatosensory cortex. The primary gustatory cortex is where the perception of a particular taste is processed.



Three nerves carry taste signals to the brain stem: the chorda tympani nerve (from the front of the tongue), the glossopharyngeal nerve (from the back of the tongue) and the vagus nerve (from the throat area and palate). In addition, the trigeminal nerve carries signals from the

touch / temperature / pain system. Taste signals combine in the brain stem areas involved in arousal (i.e. from sleep) then with smell signals in the brain to produce the sensation of flavor.