

Name : Omosebiola jesutosimi

Department : pharmacology

Matric no : 18/mhs07/043

Course : phs 212

Discuss the somatosensory system ;

The **somatosensory system** is a part of the sensory nervous system. The somatosensory system is a complex system of sensory neurons and neural pathways that responds to changes at the surface or inside the body. The axons (as afferent nerve fibers) of sensory neurons connect with, or respond to, various receptor cells. These sensory receptor cells are activated by different stimuli such as heat and nociception, giving a functional name to the responding sensory neuron, such as a thermoreceptor which carries information about temperature changes. Other types include mechanoreceptors, chemoreceptors, and nociceptors which send signals along a sensory nerve to the spinal cord where they may be processed by other sensory neurons and then relayed to the brain for further processing. Sensory receptors are found all over the body including the skin, epithelial tissues, muscles, bones and joints, internal organs, and the cardiovascular system.

The somatosensory system is distributed throughout all major parts of our body. It is responsible for sensing touch, temperature, posture, limb position, and more. It includes both sensory receptor neurons in the periphery. A somatosensory pathway will typically consist of three neurons: primary, secondary, and tertiary.

In the periphery, the primary neuron is the sensory receptor that detects sensory stimuli like touch or temperature. The cell body of the primary neuron is housed in the dorsal root ganglion of a spinal nerve or, if sensation is in the head or neck, the ganglia of the trigeminal or cranial nerves. The secondary neuron acts as a relay and is located in either the spinal cord or the brainstem. This neuron's ascending axons will cross, or decussate, to the opposite side of the spinal cord or brainstem and travel up the spinal cord to the brain, where most will terminate in either the thalamus or the cerebellum. Tertiary neurons have cell bodies in the thalamus and project to the postcentral gyrus of the parietal lobe, forming a sensory homunculus

in the case of touch. Regarding posture, the tertiary neuron is located in the cerebellum.

The primary somatosensory area of the human cortex is located in the postcentral gyrus of the parietal lobe. The postcentral gyrus is the location of the primary somatosensory area, the area of the cortex dedicated to the processing of touch information. At this location there is a map of sensory space referred to as a sensory homunculus.

A cortical homunculus is the brain's physical representation of the human body; it is a neurological map of the anatomical divisions of the body. The surface area of cortex dedicated to a body part correlates with the amount of somatosensory input from that area. For example, there is a large area of cortex devoted to sensation in the hands, while the back requires a much smaller area. Somatosensory information involved with proprioception and posture is processed in the cerebellum. The somatosensory system functions in the body's periphery, spinal cord, and the brain.