

NAME: FASIPE OLUBUNMI ASHLEY

MATRIC NO: 18/MHS02/084

DEPARTMENT: NURSING SCIENCE

COURSE TITLE: PHYSIOLOGY

COURSE CODE: 212

QUESTIONS

1. ELUCIDATE THE PATHWAY INVOLVED IN TASTE

The tongue contains small bumps called papillae, within or near which taste buds are situated. In the tongue's taste buds, the taste receptors receive sensory input via two important mechanisms – depolarization and neurotransmitter release. Intake of salty foods leads more sodium ions to enter the receptor, causing the said mechanisms. The same is true with intake of sour foods (hydrogen ions) and sweet foods (sugar molecules), both of which result to the closing of K⁺ channels upon their entry.

From the axons of the taste receptors, the sensory information is transferred to the three taste pathways via the branches of cranial nerves VII, IX and X. The chorda tympani of CN VII (facial nerve) carries the taste sensory input from the tongue's anterior two-thirds. Then, the rest of the taste sensations from the throat, palate and posterior tongue are transmitted by the branches of CN IX (glossopharyngeal nerve) and CN X (vagus nerve). From these cranial nerves, taste sensory input travels through the nerve fiber synapses to the solitary tract, the ventral posteromedial thalamic nuclei, and the thalamus. In these three locations, there are clustered neurons which respond to the same taste (sweet, sour, salty or bitter). The thalamus relays the information to the primary gustatory cortex located in the somatosensory cortex. The primary gustatory cortex is where the perception of a particular taste is processed.

The tongue contains small bumps called papillae, within or near which taste buds are situated. In the tongue's taste buds, the taste receptors receive sensory input via two important mechanisms – depolarization and neurotransmitter release. Intake of salty foods leads more sodium ions to enter the receptor, causing the said mechanisms. The same is true with intake of sour foods (hydrogen ions) and sweet foods (sugar molecules), both of which result to the closing of K⁺ channels upon their entry.

From the axons of the taste receptors, the sensory information is transferred to the three taste pathways via the branches of cranial nerves VII, IX and X. The chorda tympani of CN VII (facial nerve) carries the taste sensory input from the tongue's anterior two-thirds. Then, the rest of the taste sensations from the throat, palate and posterior tongue are transmitted by the branches of CN IX (glossopharyngeal nerve) and CN X (vagus nerve). From these cranial nerves, taste sensory input travels through the nerve fiber synapses to

the solitary tract, the ventral posteromedial thalamic nuclei, and the thalamus. In these three locations, there are clustered neurons which respond to the same taste (sweet, sour, salty or bitter). The thalamus relays the information to the primary gustatory cortex located in the somatosensory cortex. The primary gustatory cortex is where the perception of a particular taste is processed.

The sense of taste is mediated by taste receptor cells which are bundled in clusters called taste buds. Taste receptor cells sample oral concentrations of a large number of small molecules and report a sensation of taste to centers in the brainstem.

Taste buds are composed of groups of between 50 and 150 columnar taste receptor cells bundled together like a cluster of bananas. The taste receptor cells within a bud are arranged such that their tips form a small taste pore, and through this pore extend microvilli from the taste cells. The microvilli of the taste cells bear taste receptors. Interwoven among the taste cells in a taste bud is a network of dendrites of sensory nerves called "taste nerves". When taste cells are stimulated by binding of chemicals to their receptors, they depolarize and this depolarization is transmitted to the taste nerve fibers resulting in an action potential that is ultimately transmitted to the brain.

One interesting aspect of this nerve transmission is that it rapidly adapts - after the initial stimulus, a strong discharge is seen in the taste nerve fibers but within a few seconds, that response diminishes to a steady-state level of much lower amplitude.

Once taste signals are transmitted to the brain, several efferent neural pathways are activated that are important to digestive function. For example, tasting food is followed rapidly by increased salivation and by low level secretory activity in the stomach.

Among humans, there is substantial difference in taste sensitivity. Roughly one in four people is a "supertaster" that is several times more sensitive to bitter and other tastes than those that taste poorly. Such differences are heritable and reflect differences in the number of fungiform papillae and hence taste buds on the tongue.

In addition to signal transduction by taste receptor cells, it is also clear that the sense of smell profoundly affects the sensation of taste. Think about how tastes are blunted and sometimes different when your sense of smell is disrupted due to a cold.

Taste Sensations

The sense of taste is equivalent to excitation of taste receptors, and receptors for a large number of specific chemicals have been identified that contribute to the reception of taste. Despite this complexity, five types of tastes are commonly recognized by humans:

Sweet - usually indicates energy rich nutrients

Umami - the taste of amino acids (e.g. meat broth or aged cheese)

Salty - allows modulating diet for electrolyte balance

Sour - typically the taste of acids

Bitter - allows sensing of diverse natural toxins

Taste Receptors

A very large number of molecules elicit taste sensations through a rather small number of

taste receptors. Furthermore, it appears that individual taste receptor cells bear receptors for one type of taste. In other words, within a taste bud, some taste receptor cells sense sweet, while others have receptors for bitter, sour, salty and umami tastes. Much of this understanding of taste receptors has derived from behavioral studies with mice engineered to lack one or more taste receptors.

The pleasant tastes (sweet and umami) are mediated by a family of three T1R receptors that assemble in pairs. Diverse molecules that lead to a sensation of sweet bind to a receptor formed from T1R2 and T1R3 subunits. Cats have a deletion in the gene for T1R2, explaining their non-responsiveness to sweet tastes. Also, mice engineered to express the human T1R2 protein have a human-like response to different sweet tastes. The receptor formed as a complex of T1R1 and T1R3 binds L-glutamate and L-amino acids, resulting the umami taste.

The bitter taste results from binding of diverse molecules to a family of about 30 T2R receptors. Sour tasting itself involves activation of a type of TRP (transient receptor potential) channel. Surprisingly, the molecular mechanisms of salt taste reception are poorly characterized relative to the other tastes.