

NAME: MUHAMMAD-JAMIU HAMEEDAH OMOLARA

MATRICULATION NUMBER: 18/MHS02/114

DEPARTMENT: NURSING SCIENCE

COURSE TITLE: PHYSIOLOGY

COURSE CODE: PHS212

Taste is a sense that develops through the interaction of dissolved molecules with taste buds.

Currently five sub-modalities (tastes) are recognized, including sweet, salty, bitter, sour, and umami (the taste of protein). Umami is the most recent taste sensation described.

Taste is associated mainly with the tongue, although there are taste (gustatory) receptors on the palate and epiglottis as well. The surface of the tongue, along with the rest of the oral cavity, is lined by a stratified squamous epithelium. In the surface of the tongue are raised bumps, called papillae, that contain the taste buds.

There are three types of papilla, based on their appearance: vallate, foliate, and

fungi form. The number of taste buds within papillae varies, with each bud containing several specialized taste cells (gustatory receptor cells) for the transduction of taste stimuli. These receptor cells release neurotransmitters when certain chemicals in ingested substances (such as food) are carried to their surface in saliva.

Neurotransmitter from the gustatory cells can activate the sensory neurons in the facial and glossopharyngeal cranial nerves.

Primary Taste Sensations

As aforementioned, five different taste sensations are currently recognized. The first, salty, is simply the sense of Na^+ concentration in the saliva. As the Na^+ concentration becomes high outside the taste cells, a strong concentration gradient drives the diffusion into the cells. This depolarizes the cells, leading them to release neurotransmitter.

The sour taste is transduced similar to that of salty, except that it is a response to the H^+ concentration released from acidic substances (those with low pH), instead of a response to Na^+ . For example, orange juice, which contains citric acid, will taste sour because it has a pH value of about 3. Of course, it is often sweetened so that the sour taste is masked. As the concentration of the hydrogen ions increases because of ingesting acidic compounds, the depolarization of specific taste cells increases

The other three tastes; sweet, bitter and umami are transduced through G-protein coupled cell surface receptors instead of the direct diffusion of ions like we discussed with salty and sour. The sweet taste is the sensitivity of taste cells to the presence of glucose dissolved in the saliva. Molecules that are similar in structure to glucose will have a similar effect on the sensation of sweetness. Other monosaccharides such as fructose or artificial sweeteners like aspartame (Nutrasweet"), saccharine, or sucralose (Splenda") will activate the sweet receptors as well. The affinity for each of these molecules varies, and some will taste "sweeter" than glucose because they bind to the G-protein coupled receptor differently.

The bitter taste can be stimulated by a large number of molecules collectively known as alkaloids. Alkaloids are essentially the opposite of acids, they contain basic in the sense of pH) nitrogen atoms within their structures. Most alkaloids originate from plant sources, with common examples being hops (in beer), tannins (in wine), tea, aspirin, and similar molecules. Coffee contains alkaloids and is slightly acidic, with the alkaloids contributing the bitter taste to coffee. When enough alkaloids are contained in a substance it can stimulate the gag reflex. This is a protective mechanism because alkaloids are often produced by plants as a toxin to deter infectious microorganisms and plant eating animals. Such molecules may be toxic to animals as well, so we tend to avoid eating bitter foods. When we do eat bitter foods, they are often combined with a sweet component to make them more palatable (cream and sugar in coffee, for example).

The taste known as umami is often referred to as the savory taste. The name was created by the Japanese researcher who originally described it. Like sweet and bitter, it is based on the activation of G-protein coupled receptors, in this case by amino acids, especially glutamine. Thus, umami might be considered the taste of proteins, and is most associated with meat containing dishes.

Gustatory Nerve Impulses

Once the taste cells are activated by molecules liberated from the things we ingest, they release neurotransmitters onto the dendrites of sensory neurons. These neurons are part of the facial and glossopharyngeal cranial nerves, as well as a component within the vagus nerve dedicated to the gag reflex. The facial nerve connects to taste buds in the anterior third of the tongue.

The glossopharyngeal nerve connects to taste buds in the posterior two thirds of the tongue.

The vagus nerve connects to taste buds in the extreme posterior of the tongue, verging on the pharynx, which are more sensitive to noxious stimuli like bitterness.

Axons from the three cranial nerves carrying taste information travel to the medulla. From there much of the information is carried to the thalamus and then routed to the primary gustatory cortex, located near the inferior margin of the postcentral gyrus. It is the primary gustatory cortex that is responsible for our sensations of taste. And, although this region receives significant input from taste buds, it is likely that it also receives information about the smell and texture of food, all contributing to our overall taste experience. The nuclei in the medulla also send projections to the hypothalamus and amygdala, which are involved in autonomic reflexes such as gagging and salivation.