

Name: Olashina-Sadiq Bolatito

Matric number:18/MHS01/284

College/Department: MHS/MBBS

Assignment

1.MEAN ARTERIAL BLOOD PRESSURE

Mean arterial blood pressure is the average pressure existing in the arteries. It is not the arithmetic mean of systolic and diastolic pressures. It is the diastolic pressure plus one third of pulse pressure. To determine the mean pressure, diastolic pressure is considered than the systolic pressure. It is because, the diastolic period of cardiac cycle is longer (0.53 second) than the systolic period (0.27 second).

Normal mean arterial pressure: 93 mm Hg ($80 + 13 = 93$).

Formula to calculate mean arterial blood pressure:

Mean arterial blood pressure

= Diastolic pressure + 1/3 of pulse pressure

= $80 + 40/3 = 93.3$ mm Hg.

2a. PULMONARY CIRCULATION

Pulmonary circulation is the system of transportation that shunts de-oxygenated blood from the heart to the lungs to be re-saturated with oxygen before being dispersed into systemic circulation. Deoxygenated blood from the lower half of the body enters the heart from the inferior vena cava while deoxygenated blood from the upper body is delivered to the heart via the superior vena cava. Both the superior vena cava and inferior vena cava empty blood into the right atrium. Blood flows through the tricuspid valve into the right ventricle. It then flows through the pulmonic valve into the pulmonary artery before being delivered to the lungs. While in the lungs, blood diverges into the numerous pulmonary capillaries where it releases carbon dioxide and is replenished with oxygen. Once fully saturated with oxygen, the blood is transported via the pulmonary vein into the left atrium which pumps blood through the mitral valve and into the left ventricle. With a powerful contraction, the left ventricle expels oxygen-rich blood through the aortic valve and into the aorta: This is the beginning of systemic circulation.

CIRCLE OF WILLIS

The circle of Willis is a ring of interconnecting arteries located at the base of the brain around the optic chiasm or chiasma (partial crossing of the optic nerve – CNII; this crossing is important for binocular vision), infundibulum of the pituitary stalk and the hypothalamus.

This arterial ring provides blood to the brain and neighbouring structures.

Polygonal anastomotic shape offers the possibility of alternate pathways for the blood flow, which is essential for the brain functioning, since it is the structure that is mostly sensitive to hypoxia. Hypoxia of the brain tissue that lasts longer than 6 minutes results with the irreversible changes in the brain parenchyma, and

depending on the location of the lesion, the functional damages vary widely. This arterial circle is more accurately referred to as “the polygon of Willis” by the French. Although it was noticed briefly and incompletely by ancient doctors, it is described completely by an English doctor called “Thomas Willis” in his book – Cerebri Anatome in 1664. Hence the circle was named after him.

Circulus arteriosus is formed by two interconnecting arterial sources – the internal carotid arteries and the vertebro-basilar system (which is formed by two vertebral arteries and the basilar artery). This anastomosis is in an “anteroposterior order”, with the anterior source from the internal carotid arteries and their branches, and the posterior source from the vertebral arteries and their branches (specifically, from the vertebral arteries, basilar arteries and their tributaries).

SPLANCHNIC CIRCULATION

The splanchnic circulation is composed of gastric, small intestinal, colonic, pancreatic, hepatic, and splenic circulations, arranged in parallel with one another. The three major arteries that supply the splanchnic organs, celiac and superior and inferior mesenteric, give rise to smaller arteries that anastomose extensively. The circulation of some splanchnic organs is complicated by the existence of an intramural circulation. Redistribution of total blood flow between intramural vascular circuits may be as important as total blood flow. Numerous extrinsic and intrinsic factors influence the splanchnic circulation. Extrinsic factors include general hemodynamic conditions of the cardiovascular system, autonomic nervous system, and circulating neurohumoral agents. Intrinsic mechanisms include special properties of the vasculature, local metabolites, intrinsic nerves, paracrine substances, and local hormones. The existence of a multiplicity of regulatory mechanisms provides overlapping controls and restricts radical changes in tissue perfusion.

CORONARY CIRCULATION

*Coronary circulation, part of the systemic **circulatory system** that supplies blood to and provides drainage from the tissues of the **heart**. In the human heart, two coronary arteries arise from the **aorta** just beyond the semilunar valves; during **diastole**, the increased aortic pressure above the valves forces blood into the coronary arteries and thence into the musculature of the heart. Deoxygenated blood is returned to the chambers of the heart via coronary veins; most of these converge to form the coronary **venous sinus**, which drains into the right **atrium**.*

*The heart normally extracts 70 to 75 percent of the available oxygen from the blood in coronary circulation, which is much more than the amount extracted by other organs from their circulations—e.g., 40 percent by resting **skeletal muscle** and 20 percent by the liver. Obstruction of a **coronary artery**, depriving the heart tissue of oxygen-rich blood, leads to death of part of the **heart muscle** (**myocardial infarction**) in severe cases, and total **heart failure** and death may ensue.*

CUTANEOUS CIRCULATION

The cutaneous circulation is the circulation and blood supply of the skin. The skin is not a very metabolically active tissue and has relatively small energy requirements, so its blood supply is different to that of other tissues.

*Some of the circulating blood volume in the skin will flow through will flow through **arteriovenous anastomoses (AVAs)** instead of capillaries. AVAs serve a role in temperature regulation. In this article we shall consider the different adaptations of the cutaneous circulation, and its role in body temperature control.*

3. CARDIOVASCULAR ADJUSTMENT DURING EXERCISES

The integrated response to severe exercise involves fourfold to fivefold increases in cardiac output, which are due primarily to increases in cardiac rate and to a lesser extent to augmentation of stroke volume. The increase in stroke volume is partly due to an increase in end-diastolic cardiac size (Frank-Starling mechanism) and secondarily due to a reduction in end-systolic cardiac size. The full role of the Frank-Starling mechanism is masked by the concomitant tachycardia. The reduction in end-systolic dimensions can be related to increased contractility, mediated by beta adrenergic stimulation. Beta adrenergic blockade prevents the inotropic response, the decrease in end-systolic dimensions, and approximately 50% of the tachycardia of exercise. The enhanced cardiac output is distributed preferentially to the exercising muscles including the heart. Blood flow to the heart increases fourfold to fivefold as well, mainly reflecting the augmented metabolic requirements of the myocardium due to near maximal increases in cardiac rate and contractility. Blood flow to the inactive viscera (e.g., kidney and gastrointestinal tract) is maintained during severe exercise in the normal dog. It is suggested that local autoregulatory mechanisms are responsible for maintained visceral flow in the face of neural and hormonal autonomic drive, which acts to constrict renal and mesenteric vessels and to reduce blood flow. However, in the presence of circulatory impairment, where oxygen delivery to the exercising muscles is impaired as occurs to complete heart block where normal heart rate increases during exercise are prevented, or in congestive right heart failure, where normal stroke volume increases during exercise are impaired, or in the presence of severe anemia, where oxygen-carrying capacity of the blood is limited, visceral blood flows are reduced drastically and blood is diverted to the exercising musculature. Thus,, visceral flow is normally maintained during severe exercise as long as all other compensatory mechanisms remain intact. However, when any other compensatory mechanism is disrupted (even the elimination of splenic reserve in the dog), reduction and diversion of visceral flow occur.