

NAME: KELECHI AAMECHI OMEIRE

MAT NO: 18/MHS01/074

LEVEL : 200

COURSE: PHYSIOLOGY

DR OLANIYI K.S.

1. Discuss the long-term regulation of mean arterial blood pressure?

- Long-Term Regulation of Blood Pressure

There are several physiological mechanisms that regulate blood pressure in the long-term, the first of which is the renin-angiotensin-aldosterone system (RAAS).

Renin-Angiotensin-Aldosterone System (RAAS)

Renin is a peptide hormone released by the granular cells of the juxtaglomerular apparatus in the kidney. It is released in response to:

Sympathetic stimulation

Reduced sodium-chloride delivery to the distal convoluted tubule

Decreased blood flow to the kidney

Renin facilitates the conversion of angiotensinogen to angiotensin I which is then converted to angiotensin II using angiotensin-converting enzyme (ACE).

Angiotensin II is a potent vasoconstrictor. It acts directly on the kidney to increase sodium reabsorption in the proximal convoluted tubule. Sodium is reabsorbed via the sodium-hydrogen exchanger. Angiotensin II also promotes release of aldosterone.

ACE also breaks down a substance called bradykinin which is a potent vasodilator. Therefore, the breakdown of bradykinin potentiates the overall constricting effect.

Aldosterone promotes salt and water retention by acting at the distal convoluted tubule to increase expression of epithelial sodium channels. Furthermore, aldosterone increases the activity of the basolateral sodium-potassium ATP-ase, thus increasing the electrochemical gradient for movement of sodium ions.

More sodium collects in the kidney tissue and water then follows by osmosis. This results in decreased water excretion and therefore increased blood volume and thus blood pressure.

Anti-Diuretic Hormone (ADH)

The second mechanism by which blood pressure is regulated is release of Anti Diuretic Hormone (ADH) from the OVLT of the hypothalamus in response to thirst or an increased plasma osmolarity.

ADH acts to increase the permeability of the collecting duct to water by inserting aquaporin channels (AQP2) into the apical membrane.

It also stimulates sodium reabsorption from the thick ascending limb of the loop of Henle. This increases water reabsorption thus increasing plasma volume and decreasing osmolarity.

2 Write short notes on the following:

- **Pulmonary circulation**, is a system of **blood** vessels that forms a closed circuit between the heart and the **lungs**, as distinguished from the systemic circulation between the **heart** and all other body tissues. Deoxygenated blood from the lower half of the body enters the heart from the inferior vena cava while deoxygenated blood from the upper body is delivered to the heart via the superior vena cava. Both the superior vena cava and inferior vena cava empty blood into the right atrium. Blood flows through the tricuspid valve into the right ventricle. It then flows through the pulmonic valve into the pulmonary artery before being delivered to the

lungs. While in the lungs, blood diverges into the numerous pulmonary capillaries where it releases carbon dioxide and is replenished with oxygen. Once fully saturated with oxygen, the blood is transported via the pulmonary vein into the left atrium which pumps blood through the mitral valve and into the left ventricle. With a powerful contraction, the left ventricle expels oxygen-rich blood through the aortic valve and into the aorta: This is the beginning of systemic circulation.

- **The circle of Willis** is a ring of interconnecting arteries located at the base of the brain around the optic chiasm or chiasma (partial crossing of the **optic nerve**– CNII; this crossing is important for binocular vision), infundibulum of the pituitary stalk and the **hypothalamus**. This arterial ring provides blood to the brain and neighboring structures.
- **The splanchnic circulation** is composed of gastric, small intestinal, colonic, pancreatic, hepatic, and splenic circulations, arranged in parallel with one another. The three major arteries that supply the splanchnic organs, celiac and superior and inferior mesenteric, give rise to smaller arteries that anastomose extensively. The circulation of some splanchnic organs is complicated by the existence of an intramural circulation. Redistribution of total blood flow between intramural vascular circuits may be as important as total blood flow. Numerous extrinsic and intrinsic factors influence the splanchnic circulation. Extrinsic factors include general hemodynamic conditions of the cardiovascular system, autonomic nervous system, and circulating neurohumoral agents. Intrinsic mechanisms include special properties of the vasculature, local metabolites, intrinsic nerves, paracrine substances, and local hormones. The existence of a multiplicity of regulatory mechanisms provides overlapping controls and restricts radical changes in tissue perfusion.
- **Coronary circulation** is the **circulation of blood** in the **blood vessels** that supply the **heart muscle** (myocardium). **Coronary arteries** supply **oxygenated** blood to the heart muscle, and **cardiac veins** drain away the blood once it has been deoxygenated. Because the rest of the body, and most especially the **brain**, needs a steady supply of oxygenated blood that is free of all but the slightest interruptions, the

heart is required to function continuously. Making it a circulation of major importance, not only to its tissue but to the entire body and even the level of consciousness of the brain. Any interruptions of coronary circulation quickly causes heart attack.

- **The cutaneous** circulation is the circulation and blood supply of the skin. The skin is not a very metabolically active tissue and has relatively small energy requirements, so its blood supply is different to that of other tissues. Some of the circulating blood volume in the skin will flow through will flow through **arteriovenous anastomoses (AVAs)** instead of capillaries. AVAs serve a role in temperature regulation.

3 Discuss the cardiovascular adjustment that occurs during exercise?

- The integrated response to severe exercise involves fourfold to fivefold increases in cardiac output, which are due primarily to increases in cardiac rate and to a lesser extent to augmentation of stroke volume. The increase in stroke volume is partly due to an increase in end-diastolic cardiac size (Frank-Starling mechanism) and secondarily due to a reduction in end-systolic cardiac size. The full role of the Frank-Starling mechanism is masked by the concomitant tachycardia. The reduction in end-systolic dimensions can be related to increased contractility, mediated by beta adrenergic stimulation. Beta adrenergic blockade prevents the inotropic response, the decrease in end-systolic dimensions, and approximately 50% of the tachycardia of exercise. The enhanced cardiac output is distributed preferentially to the exercising muscles including the heart. Blood flow to the heart increases fourfold to fivefold as well, mainly reflecting the augmented metabolic requirements of the myocardium due to near maximal increases in cardiac rate and contractility. Blood flow to the inactive viscera (e.g., kidney and gastrointestinal tract) is maintained during severe exercise in the normal dog. It is suggested that local autoregulatory mechanisms are responsible for maintained visceral flow in the face of neural and hormonal autonomic drive, which acts to constrict

renal and mesenteric vessels and to reduce blood flow. However, in the presence of circulatory impairment, where oxygen delivery to the exercising muscles is impaired as occurs to complete heart block where normal heart rate increases during exercise are prevented, or in congestive right heart failure, where normal stroke volume increases during exercise are impaired, or in the presence of severe anemia, where oxygen-carrying capacity of the blood is limited, visceral blood flows are reduced drastically and blood is diverted to the exercising musculature. Thus, visceral flow is normally maintained during severe exercise as long as all other compensatory mechanisms remain intact. However, when any other compensatory mechanism is disrupted (even the elimination of splenic reserve in the dog), reduction and diversion of visceral flow occur.