

NAME :OLAWUYI RAODAT FEYISEIKEMI

MATRIC NO: 17/MHS01/255

ASSIGNMENT TITLE: Renal physiology for mbbs student

COURSE TITLE: Renal physiology body fluid and temperature regulation

COURSE CODE: phs 303

QUESTION

1 Discuss the role of kidney in glucose homeostasis

ANSWER

They release glucose into the circulation via gluconeogenesis, uptake of glucose from the circulation to satisfy their energy needs, and reabsorption of glucose at the level of the proximal tubule. Renal release of glucose into the circulation is the result of glycogenolysis and gluconeogenesis, respectively involving the breaking down and formation of glucose-6-phosphate from precursors (eg, lactate, glycerol, amino acids). With regard to renal reabsorption of glucose, the kidneys normally retrieve as much glucose as possible, rendering the urine virtually glucose free. The glomeruli filter from plasma approximately 180 grams of D-glucose per day, all of which is reabsorbed through glucose transporter proteins that are present in cell membranes within the proximal tubules. If the capacity of these transporters is exceeded, glucose appears in the urine. The process of renal glucose reabsorption is mediated by active (sodium-coupled glucose cotransporters) and passive (glucose transporters) transporters. In hyperglycemia, the kidneys may play an exacerbating role by reabsorbing excess glucose, ultimately contributing to chronic hyperglycemia, which in turn contributes to chronic glycemic burden and the risk of microvascular consequences. This article provides an extensive review of the kidneys role in normal human physiology, the mechanisms by which they contribute to glucose regulation, and the potential impact of glucose imbalance on the kidneys.

2 Discuss the process of micturition

ANSWER

Micturition Process

Micturition process consists of two phases:

- Storage phase
- Voiding phase

Storage Phase

The urinary bladder is a balloon-shaped, hollow, muscular, organ that acts as the storage organ for urine. The urinary bladder in a healthy urinary system can store up to 16 ounces of urine for 2 to 5 hours easily. The circular sphincter muscles prevent leakage of urine. They close tightly around the opening of the bladder into the tube (urethra) that allows the passage of urine outside the body.

Voiding Phase

When the bladder is filled with urine, the nerves in it are triggered, which in turn stimulates the need to urinate. The brain signals urinary bladder to contract. The receptors of the urinary bladder send a signal to the central nervous system, in response to which the nervous system sends a signal that incites the contraction of the urinary bladder. Through the urinary opening at the urethra, the urine is eliminated, and the process is called micturition. The neural mechanism involved is called the micturition reflex.

3 Explain juxtaglomerular apparatus

ANSWER

Juxtaglomerular apparatus is a specialized organ situated near the glomerulus of each nephron (juxta = near).

STRUCTURE OF JUXTAGLOMERULAR APPARATUS

Juxtaglomerular apparatus is formed by three different structures:

1. Macula densa
2. Extraglomerular mesangial cells
3. Juxtaglomerular cells

MACULA Densa: Macula densa is the end portion of thick ascending segment before it opens into distal convoluted tubule. It is situated between afferent and efferent arterioles of the same nephron. It is very close to afferent arteriole. Macula densa is formed by tightly packed cuboidal epithelial cells.

EXTRAGLOMERULAR MESANGIAL CELLS: Extraglomerular mesangial cells are situated in the triangular region bound by afferent arteriole, efferent arteriole and macula densa. These cells are also called **agranular cells, lacis cells** or

Goormaghtigh cells.

JUXTAGLOMERULAR CELLS: Juxtaglomerular cells are specialized smooth muscle cells situated in the wall of afferent arteriole just before it enters the Bowman capsule. These smooth muscle cells are mostly present in tunica media and tunica adventitia of the wall of the afferent arteriole.

Juxtaglomerular cells are also called **granular cells** because of the presence of secretory granules in their cytoplasm.

FUNCTIONS OF JUXTAGLOMERULAR APPARATUS

Primary function of juxtaglomerular apparatus is the secretion of hormones. It also regulates the glomerular blood flow and glomerular filtration.

4 Discuss the role of kidney in regulation of blood pressure

ANSWER

Kidneys play an important role in the long-term regulation of arterial blood pressure by two ways:

- By regulating the volume of extracellular fluid
- through renin-angiotensin mechanism.

5 Discuss the role of kidney in calcium homeostasis

ANSWER

kidney plays a key role in this process by the fine regulation of calcium excretion. More than 95% of filtered calcium is reabsorbed along the renal tubules. In the proximal tubules, 60% of filtered calcium is reabsorbed by passive mechanisms. In the thick ascending limb, 15% of calcium is reabsorbed by paracellular diffusion through paracellin-1 (claudin-16). The calcium sensing receptor (CaSR) in the basolateral membrane of the thick ascending limb senses the change in iCa^{2+} and inhibits calcium reabsorption independent to PTH and $1,25(OH)_2D_3$. The fine regulation of calcium excretion occurs in the distal convoluted tubules and connecting tubules despite the fact that only 10-15% of filtered calcium is reabsorbed there. Transient receptor potential vanilloid 5 (TRPV5) and 6 (TRPV6) in the apical membrane act as the main portal of entry, calbindin-D28K delivers Ca^{2+} in the cytoplasm, and then Na^+/Ca^{2+} exchanger (NCX1) and plasma membrane Ca^{2+} -ATPase in the basolateral membrane serve as an exit. In the cortical collecting duct, TRPV6 is expressed, but the role might be negligible. In addition to PTH and $1,25(OH)_2D_3$, acid-base disturbance, diuretics, and estrogen affect on these calcium channels. Recently, klotho and fibroblast growth factor 23 (FGF23) are suggested as new players in the calcium metabolism. Klotho is exclusively expressed in the kidney and co-localized with TRPV5, NCX1, and calbindin-D28K. Klotho increases calcium reabsorption through trafficking of TRPV5 to the plasma membrane, and also converts FGF receptor to the specific FGF23 receptor. FGF23:klotho complex bound to FGF receptor inhibits 1α -hydroxylase of vitamin D, and contributes to calcium reabsorption and phosphate excretion in the kidney.