

18/MHS07/044

PHS 212

Assignment: write short note on micturition

Answer:

## **Micturition**

Micturition is also known as the **voiding** phase of bladder control and it is typically a short-lasting event. Urinary flow rate in a full bladder is:

20-25ml/s in men

25-30ml/s in women

Whilst the capacity of the bladder varies from roughly 300-550ml, afferent nerves in the bladder wall signal the need to void the bladder at around **400ml** of filling.

### **Regulation of Micturition**

Passing of urine is under **parasympathetic** control. Bladder afferents signals ascend through the spinal cord and then project to the pontine micturition centre and cerebrum. Upon the voluntary decision to urinate, neurones of the pontine micturition centre fire to excite the sacral preganglionic neurones.

There is subsequent parasympathetic stimulation to the **Pelvic Nerve (S2-4)** causing a release of ACh, which works on M3 muscarinic ACh receptors on the detrusor muscle, causing it to contract and increase intra-vesicular pressure. The pontine micturition centre also inhibits Onuf's nucleus, with a resultant reduction in sympathetic stimulation to the internal urethral sphincter causing relaxation.

Finally, a **conscious reduction** in voluntary contraction of the external urethral sphincter from the cerebral cortex allows for distention of the urethra and the passing of urine. In the female, urination is assisted by gravity, while in the male, bulbospongiosus contractions and squeezing along the length of the penis helps to expel all of the urine.

## **Stages of Micturition**

The urinary bladder has two distinct stages or phases:

- 1 Resting or filling stage
- 2 Voiding stage

### **Resting or Filling Stage**

It is in this phase of the bladder that the urine is transported from the kidneys via the ureters into the bladder. The ureters are thin muscular tubes that arise from each of the kidneys and extend downwards where they enter the bladder obliquely.

The oblique placement of the ureters in the bladder wall serves a very important function. The opening of the ureter into the urinary bladder is not guarded by any sphincter or muscle. Therefore, this oblique nature of opening prevents the urine from re-entering the ureters. At the same time, the main muscle of the urinary bladder, the detrusor muscle, is relaxing allowing the bladder to distend and accommodate more urine.

### **Voiding Stage**

During this stage, both the urinary bladder and the urethra come into play together. The detrusor muscle of the urinary bladder which was relaxing so far starts to contract once the bladder's storage capacity is reached.

The urethra is controlled by two sets of muscles: The internal and external urethral sphincters. The internal sphincter is a smooth muscle whereas the external one is skeletal. Both these sphincters are in a contracted state during the filling stage.

As mentioned earlier, the process of micturition is governed by both the nervous and muscular systems. Within the nervous system, the process is governed by the autonomous nervous system and the somatic system. Once the urinary bladder reaches its maximum capacity, the stretch receptors in the walls of the bladder send an impulse via the pelvic nerve to the brain via the spinal cord.

The micturition reflex is ultimately generated from the level of the spinal cord after it receives reflexes from the pontine region in the brain. Once the bladder and the urethra receive the signals to empty the bladder, the two sphincters relax and the detrusor muscle causes the contractions of the bladder.

Along with these muscles, the muscles of the abdomen also play a role by putting pressure on the bladder wall. This leads to complete emptying of the bladder.