

**NAME; EYEREGBA LOIS EJIROGHENE**

**DEPARTMENT; NURSINCE SCIENCE**

**LEVEL; 200**

**MATRIC NO 18/MHS02/077**

**COURSE; PHS 212**

**QUESTION**

**WRITE A SHORT NOTE ON MICTURITION**

Micturition or urination is the process of expelling urine from the bladder. This act is also known as voiding of the bladder. The excretory system in humans includes a pair of kidneys, two ureters, a urinary bladder and a urethra. The kidneys filter the urine and it is transported to the urinary bladder via the ureters where it is stored till its expulsion. The process of micturition is regulated by the nervous system and the muscles of the bladder and urethra. The urinary bladder can store around 350-400ml of urine before it expels it out.

**Stages of Micturition**

The urinary bladder has two distinct stages or phases:

1. Resting or filling stage
2. Voiding stage

### **Resting or Filling Stage**

It is in this phase of the bladder that the urine is transported from the kidneys via the ureters into the bladder. The ureters are thin muscular tubes that arise from each of the kidneys and extend downwards where they enter the bladder obliquely.

The oblique placement of the ureters in the bladder wall serves a very important function. The opening of the ureter into the urinary bladder is not guarded by any sphincter or muscle. Therefore, this oblique nature of opening prevents the urine from re-entering the ureters. At the same time, the main muscle of the urinary bladder, the detrusor muscle, is relaxing allowing the bladder to distend and accommodate more urine.

### **Voiding Stage**

During this stage, both the urinary bladder and the urethra come into play together. The detrusor muscle of the urinary bladder which was relaxing so far starts to contract once the bladder's storage capacity is reached.

The urethra is controlled by two sets of muscles: **The internal and external urethral sphincters**. The internal sphincter is a smooth muscle whereas the external one is skeletal. Both these sphincters are in a contracted state during the filling stage.

### **Control of micturition**

Children and adults have considerable control over when and where they pass urine. They can also increase or decrease the rate of flow and even stop and start again, so micturition is clearly more than just a simple reflex. This control is learnt in infancy and involves other sensory fibres in the bladder wall. These fibres convey information on the degree of bladder fullness via the spine to the higher centres of the brain, the thalamus and cerebral cortex. This causes us to become aware that we need to pass urine and of the urgency of the situation.

### **Conclusion of Micturition**

As mentioned earlier, the process of micturition is governed by both the nervous and muscular systems. Within the nervous system, the process is governed by the autonomous nervous system and the somatic system. Once the urinary bladder reaches its maximum capacity, the stretch receptors in the walls of the bladder send an impulse via the pelvic nerve to the brain via the spinal cord.

The micturition reflex is ultimately generated from the level of the spinal cord after it receives reflexes from the pontine region in the brain. Once the bladder and the urethra receive the signals to empty the bladder, the two sphincters relax and the detrusor muscle causes the contractions of the bladder.

Along with these muscles, the muscles of the abdomen also play a role by putting pressure on the bladder wall. This leads to complete emptying of the bladder.